

New Proofs for Inequalities of Power-Exponential Functions

This is the Published version of the following publication

Guo, Bai-Ni and Qi, Feng (2000) New Proofs for Inequalities of Power-Exponential Functions. RGMIA research report collection, 4 (1).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository https://vuir.vu.edu.au/17378/

NEW PROOFS FOR INEQUALITIES OF POWER-EXPONENTIAL **FUNCTIONS**

BAI-NI GUO AND FENG QI

ABSTRACT. Some new proofs for the following inequalities of power-exponential functions are given

$$\frac{y^{x^y}}{x^{y^x}}>\frac{y}{x}>\frac{y^x}{x^y},\quad \left(\frac{y}{x}\right)^{xy}>\frac{y^y}{x^x},$$
 where $y>x>0$ and $(x-1)(y-1)>0.$

1. Introduction

Using different techniques, Professor Dr. F. Qi and L. Debnath proved the following in [6]:

Theorem 1. For y > x > 0 and (x - 1)(y - 1) > 0, we have

$$\frac{y^{x^y}}{x^{y^x}} > \frac{y}{x} > \frac{y^x}{x^y},\tag{1}$$

$$\left(\frac{y}{x}\right)^{xy} > \frac{y^y}{x^x}.\tag{2}$$

For 0 < x < 1 < y, the right hand sides of (1) and (2) are reversed.

If 0 < x < 1 < y or 0 < x < y < e, then

$$\frac{y^x}{x^y} > \frac{y \ln x}{x \ln y} \cdot \frac{y^x - 1}{x^y - 1} > 1.$$
 (3)

If e < x < y, then inequality (3) is reversed.

The following lemma is well-known:

Lemma 1 ([3, page 11]). If f(t) is an increasing integrable function on I, then the arithmetic mean of the function f(t),

$$\Phi(r,s) = \begin{cases} \frac{1}{s-r} \int_r^s f(t) dt, & r \neq s, \\ f(r), & r = s, \end{cases}$$
(4)

is also increasing with both r and s on I.

Date: December 26, 2000.

²⁰⁰⁰ Mathematics Subject Classification. Primary 26D07, 26D20.

Key words and phrases. Inequality, power-exponential function, arithmetic mean, monotonicity.

The author was supported in part by NSF of Henan Province (no. 004051800), SF for Pure Research of the Education Department of Henan Province (no. 1999110004), Doctor Fund of Jiaozuo Institute of Technology, and NNSF (no. 10001016) of China .

This paper was typeset using AMS-IATEX.

The above lemma was reproved and used successfully to verify monotonicity, logarithmic convexity, and other properties of the extended mean values, generalized weighted mean values, and generalized abstracted mean values in [4, 5] and [7, 8].

In this paper, using Lemma 1 and other techniques, we will give some new proofs for the inequalities of power-exponential functions presented in Theorem 1.

2. New proofs

2.1. New proof of the right hand side of inequality (1).

2.1.1. First proof. Using standard arguments, the right hand side of inequality (1) can be rewritten as

$$(1-x)\ln y > (1-y)\ln x \tag{5}$$

for y > x > 0 and (x - 1)(y - 1) > 0. Inequality (5) is equivalent to

$$\frac{\ln y}{1-y} > \frac{\ln x}{1-x}.\tag{6}$$

By Lemma 1, we may deduce that the function ϕ defined by

$$\phi(u) = \frac{\ln u}{u - 1} = \frac{1}{u - 1} \int_{1}^{u} \frac{1}{t} dt,$$

$$\phi(1) = 1$$
(7)

is decreasing for u > 0. Therefore, inequality (6) holds and the right hand side of inequality (1) follows.

2.1.2. Second proof. We can rewrite the right hand side of inequality (1) as follows.

$$y^{\frac{1}{y-1}} < x^{\frac{1}{x-1}} \tag{8}$$

for y > x > 0 and (x-1)(y-1) > 0. This is equivalent to the function $t^{\frac{1}{t-1}}$, and then the function $\frac{\ln t}{t-1}$, decreasing in t > 0. This was done in [6].

2.2. New proofs of inequality (2).

2.2.1. First proof. It is easy to see that inequality (2) is equivalent to

$$\frac{y\ln y}{y-1} > \frac{x\ln x}{x-1} \tag{9}$$

for y > x > 0 and (x - 1)(y - 1) > 0.

From Lemma 1, we obtain that the function

$$\varphi(u) = \frac{u \ln u}{u - 1} = \frac{1}{u - 1} \int_{1}^{u} (1 + \ln t) dt,$$

$$\varphi(1) = 1$$
(10)

is increasing for u > 0. Therefore, inequality (9) holds and (2) follows.

2.2.2. Second Proof. Inequality (2) can be rewritten as

$$\frac{(y^y)^x}{(x^x)^y} > \frac{y^y}{x^x},\tag{11}$$

which is equivalent to

$$y^{\frac{y}{y-1}} > x^{\frac{x}{x-1}} \tag{12}$$

for y > x > 0 and (x-1)(y-1) > 0. It suffices to prove that the function $\frac{t \ln t}{t-1}$ is increasing in t > 0. This was done in [6].

2.3. New proof of the left hand side of inequality (1). The equivalent form of the left hand side of inequality (1) is

$$\frac{x^y - 1}{y^x - 1} > \frac{\ln x}{\ln y} \tag{13}$$

for (x-1)(y-1) > 0 and y > x > 0.

Let $\omega(t) = x^{yt}$ and $\psi(t) = y^{xt}$ for $t \in [0,1]$. Direct computation yields

$$\omega'(t) = x^{yt} \cdot y \ln x,$$

$$\psi'(t) = y^{xt} \cdot x \ln y.$$
(14)

Then, using the Cauchy's mean-value theorem for differentiation and the right inequality in (1), there exists a point $\xi \in (0,1)$, such that

$$\frac{x^{y} - 1}{y^{x} - 1} = \frac{\omega(1) - \omega(0)}{\psi(1) - \psi(0)}$$

$$= \frac{\omega'(\xi)}{\psi'(\xi)}$$

$$= \left(\frac{x^{y}}{y^{x}}\right)^{\xi} \cdot \frac{y \ln x}{x \ln y}$$

$$> \left(\frac{x}{y}\right)^{\xi} \cdot \frac{y \ln x}{x \ln y}$$

$$> \left(\frac{y}{x}\right)^{1-\xi} \cdot \frac{\ln x}{\ln y}$$

$$> \frac{\ln x}{\ln y}.$$
(15)

2.4. New proofs of inequality (3). Since the inequality $x^y < y^x$ holds for 0 < x < y < e and is reversed for e < x < y, (For more information about this inequality, please refer to [1, 2, 3, 6]), from the first three lines in inequality (15), we have that

$$\frac{x^y - 1}{y^x - 1} = \left(\frac{x^y}{y^x}\right)^{\xi} \cdot \frac{y \ln x}{x \ln y} < \frac{y \ln x}{x \ln y} \tag{16}$$

holds for 0 < x < y < e and is reversed for e < x < y. This implies that the right hand side of inequality (3) holds.

In [6], it was proved that the function $\frac{t-1}{\ln t}$ is increasing in t > 0. For y > 1 > x > 0, we have $y^x > 1 > x^y$, and thus

$$\frac{y^x - 1}{\ln y^x} > \frac{x^y - 1}{\ln x^y}. (17)$$

This implies that the right hand side of inequality (3) holds.

For 0 < x < y < e, we have $y^x > x^y$, and applying the same procedure produces an identical result.

It was also proved in [6] that the function $\frac{t-1}{t \ln t}$ decreases in t>0. Since the inequality $x^y < y^x$ holds for 0 < x < 1 < y and for 0 < x < y < e, then

$$\frac{y^x - 1}{y^x \ln y^x} < \frac{x^y - 1}{x^y \ln x^y}. (18)$$

Simplification of inequality (18) gives us the left hand side of inequality (3). Similar arguments enable us to establish the reversed inequalities.

References

- P. S. Bullen, A Dictionary of Inequalities, Pitman Monographs and Surveys in Pure and Applied Mathematics 97, Addison Wesley Longman Limited, 1998.
- 2. J.-C. Kuang, Applied Inequalities (Changyong Budengshi), 2nd edition, Hunan Education Press, Changsha, China, 1993. (Chinese)
- 3. D. S. Mitrinović, Analytic Inequalities (Jiexi Budengshi), Chinese Edition, translated by X.-P. Zhang and L. Wang, Science Press, Beijing, 1987. (Chinese)
- 4. F. Qi, Generalized abstracted mean values, J. Ineq. Pure Appl. Math. 1 (2000), no. 1, Article 4. http://jipam.vu.edu.au/v1n1/013_99.html. RGMIA Res. Rep. Coll. 2 (1999), no. 5, Article 4, 633–642. http://rgmia.vu.edu.au/v2n5.html.
- 5. F. Qi, Logarithmic convexity of extended mean values, Proc. Amer. Math. Soc. (2001), in the press. RGMIA Res. Rep. Coll. **2** (1999), no. 5, Article 5, 643–652. http://rgmia.vu.edu.au/v2n5.html.
- 6. F. Qi and L. Debnath, *Inequalities of power-exponential functions*, J. Ineq. Pure Appl. Math. 1 (2000), no. 2, Article 15. http://jipam.vu.edu.au/v1n2/011_99.html.
- F. Qi, J.-Q. Mei, D.-F. Xia, and S.-L. Xu, New proofs of weighted power mean inequalities and monotonicity for generalized weighted mean values, Math. Ineq. Appl. 3 (2000), no. 3, 377–383.
- F. Qi, S.-L. Xu, and L. Debnath, A new proof of monotonicity for extended mean values, Intern. J. Math. Math. Sci. 22 (1999), no. 2, 415–420.

Department of Mathematics, Jiaozuo Institute of Technology, Jiaozuo City, Henan 454000, The People's Republic of China

Department of Mathematics, Jiaozuo Institute of Technology, Jiaozuo City, Henan 454000, The People's Republic of China

 $E ext{-}mail\ address: qifeng@jzit.edu.cn}$

URL: http://rgmia.vu.edu.au/qi.html or http://rgmia.vu.edu.au/authors/FQi.htm