

Approximation of pn by Hn

This is the Published version of the following publication

Alikhani, Akram and Hassani, Mehdi (2005) Approximation of pn by Hn. Research report collection, 8 (4).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository https://vuir.vu.edu.au/18117/

APPROXIMATION OF p_n BY H_n

AKRAM ALIKHANI MEHDI HASSANI

ABSTRACT. In this paper we introduce some bounds including $H_n = \sum_{k=1}^n \frac{1}{k}$, for p_n , n^{th} prime number. Then we observe that the Prime Number Theorem is equivalent with $p_n \sim nH_n$, when n tends to infinity.

1. Introduction

As usual, let p_n be the n^{th} prime. According to the Prime Number Theorem (PNT) [3], we know that:

$$(1.1) p_n = n \log n + o(n \log n) (n \to \infty).$$

Also, we know that [1], if $H_n = \sum_{k=1}^n \frac{1}{k}$, then:

(1.2)
$$H_n = \log n + O(1) \qquad (n \to \infty).$$

So, considering (1.1) and (1.2), we obtain:

$$p_n = n(H_n + O(1)) + o(n\log n) = nH_n + o(n\log n) \qquad (n \to \infty).$$

Therefore, comparing p_n and nH_n seems to be a nice problem. To consider this problem, we need some bounds concerning p_n and H_n , which we recall them from literatures. About p_n , we have the following bounds [4]:

$$n\log n + n\log_2 n - n + n\frac{\log_2 n - 2.25}{\log n} \le p_n \le n\log n + n\log_2 n - n + n\frac{\log_2 n - 1.8}{\log n},$$

which left hand side of it holds true for every $n \ge 2$ and the right hand side of it holds true for every $n \ge 27076$, $\log_2 n$ means $\log \log n$ and base of all logarithms is e. Also, for H_n we have the following bounds [2]:

$$(1.4) \gamma + \log(n+0.5) < H_n < \gamma + \log(n-1+e^{1-\gamma}) (n > 1),$$

where γ is Euler constant. In this note, we search some bounds of the form $nH_n + e(n)$, which we will find e(n) in both cases lower and upper bounds.

2. Inequalities of the form nH_n

Consider the following inequality:

$$(2.1) nH_n + a(n) \le p_n \le nH_n + b(n).$$

Here, we try to find some suitable functions a(n) and b(n), such that (2.1) holds true.

 $^{1991\} Mathematics\ Subject\ Classification.\ 11A41,\ 26D15.$

Key words and phrases. Primes, Harmonic series.

Upper Bound. To find above mentioned upper bound, considering (1.4), we have:

(2.2)
$$n \left(\gamma + \log(n + 0.5) \right) < nH_n \le n \left(\gamma + \log(n - 1 + e^{1 - \gamma}) \right).$$

Using left hand side of above inequality, we have:

$$n\left(\gamma + \log(n + 0.5)\right) + b(n) \le nH_n + b(n),$$

and considering this inequality, with the right hand side of (1.3), we must have:

$$n\log n + n\log_2 n - n + n\frac{\log_2 n - 1.8}{\log n} \le n\left(\gamma + \log(n + 0.5)\right) + b(n),$$

or equivalently,

$$n \log n + n \log_2 n - n + n \frac{\log_2 n - 1.8}{\log n} - n \left(\gamma + \log(n + 0.5) \right) \le b(n),$$

Since, b(n) is going to appears in upper bound for p_n , the best possible case is:

(2.3)
$$b(n) = n \log_2 n - n(1+\gamma) + n \left(\frac{\log_2 n - 1.8}{\log n} - \log \left(1 + \frac{0.5}{n} \right) \right).$$

Thus, we have:

$$(2.4) p_n \le nH_n + n\log_2 n - n(1+\gamma) + n\left(\frac{\log_2 n - 1.8}{\log n} - \log\left(1 + \frac{0.5}{n}\right)\right),$$

which holds for $n \geq 27076$.

Lower Bound. To find above mentioned lower bound, considering (2.2), we have:

$$nH_n + a(n) \le n\gamma + n\log(n - 1 + e^{1-\gamma}) + a(n).$$

Considering this inequality with the left hand side of (2.1) and the left hand side of (1.3), we must have:

$$n\gamma + n\log(n - 1 + e^{1-\gamma}) + a(n) \le n\log n + n\log_2 n - n + n\frac{\log_2 n - 2.25}{\log n}$$
.

Since, we want to find the maximum lower bound in the left hand side of (2.1), the best possible choice for a(n), is:

$$a(n) = n \log_2 n - n(1+\gamma) + n \left(\frac{\log_2 n - 2.25}{\log n} - \log \left(1 + \frac{e^{1-\gamma} - 1}{n} \right) \right).$$

So, we have:

$$(2.5) nH_n + n\log_2 n - n(1+\gamma) + n\left(\frac{\log_2 n - 2.25}{\log n} - \log\left(1 + \frac{e^{1-\gamma} - 1}{n}\right)\right) \le p_n,$$

which holds for $n \geq 2$. Therefore, considering (2.4) and (2.5), for every $n \geq 27076$, we obtain:

$$(2.6) |p_n - (nH_n + n\log_2 n - n(1+\gamma))| \le n\left(\frac{\log_2 n - 1.8}{\log n} - \log\left(1 + \frac{0.5}{n}\right)\right).$$

3. An Equivalent for the PNT

Considering (2.6), we obtain:

$$p_n = nH_n + n\log_2 n - n(1+\gamma) + O\left(\frac{n\log_2 n}{\log n}\right),$$

which is a very strong form of an equivalent for the PNT. In fact we observe that the PNT holds if and only if $p_n \sim nH_n$, when n tends to infinity. To see this, first suppose that the PNT holds true. So, when n tends to infinity, we have:

$$p_n = n \log n + o(n \log n).$$

Considering this with $H_n \sim \log n$, we obtain:

$$p_n = n(H_n + O(1)) + o(n \log n)$$

$$= nH_n + O(n) + o(n \log n)$$

$$= nH_n + o(n \log n)$$

$$= nH_n + o(n(H_n + O(1)))$$

$$= nH_n + o(nH_n) + o(O(n))$$

$$= nH_n + o(nH_n).$$

Inversely, suppose $p_n \sim nH_n$, then:

$$p_n = nH_n + o(nH_n),$$

which considering this with $H_n \sim \log n$, we obtain:

$$p_n = n(\log n + O(1)) + o(n(\log n + O(1)))$$

$$= n \log n + O(n) + o(n \log n + O(n))$$

$$= n \log n + O(n) + o(n \log n)$$

$$= n \log n + o(n \log n),$$

and this is PNT.

References

- [1] M. Abramowitz and I.A. Stegun, *HANDBOOK OF MATHEMATICAL FUNCTIONS: with Formulas, Graphs, and Mthematical Tables*, Dover Publications, 1972.
- [2] N. Batir, "Some new inequalities for Gamma and polygamma functions", RGMIA Research Report Collection, 7(3), Article 1, 2004.
- [3] H. Davenport, Multiplicative Number Theory (Second Edition), Springer-Verlag, 1980.
- [4] P. Dusart, Inégalités explicites pour $\psi(X)$, $\theta(X)$, $\pi(X)$ et les nombres premiers, C. R. Math. Acad. Sci. Soc. R. Can. 21 (1999), no. 2, 53–59.

 $\label{thm:condition} \text{Institute for Advanced, Studies in Basic Sciences, P.O. Box 45195-1159, Zanjan, Iran. } \\ E-mail\ address: \texttt{alikhani@iasbs.ac.ir, mmhassany@srttu.edu} \\$