

Inner Product Inequalities for Two Equivalent Norms and Applications

This is the Published version of the following publication

Dragomir, Sever S (2008) Inner Product Inequalities for Two Equivalent Norms and Applications. Research report collection, 11 (Supp).

The publisher's official version can be found at

Note that access to this version may require subscription.

Downloaded from VU Research Repository https://vuir.vu.edu.au/18382/

INNER PRODUCT INEQUALITIES FOR TWO EQUIVALENT NORMS AND APPLICATIONS

S. S. DRAGOMIR

ABSTRACT. Some inequalities for two inner products $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$ which generate the equivalent norms $\|\cdot\|_1$ and $\|\cdot\|_2$ with applications for invertible bounded linear operators, positive definite self-adjoint operators, integral and discrete inequalities are given.

1. INTRODUCTION

Let $(H, \langle \cdot, \cdot \rangle)$ be an inner product over the real or complex number field K. The following inequality

(1.1)
$$|\langle x, y \rangle| \le ||x|| ||y||, \qquad x, y \in H$$

is well known in the literature as *Schwarz's inequality*. It plays an essential role in obtaining various results in the Geometry of Inner Product Spaces as well as in its applications in Operator Theory, Approximation Theory and other fields.

Due to the fact that

(1.2)
$$|\operatorname{Re}\langle x, y\rangle| \le ||x|| ||y||, \quad x, y \in H_{2}$$

we can introduce the angle between the vectors x, y, denoted by $\Phi_{x,y}$, through the formula

(1.3)
$$\cos \Phi_{xy} := \frac{\operatorname{Re} \langle x, y \rangle}{\|x\| \|y\|}, \qquad x, y \neq 0.$$

As observed by Krein in 1969, [6] (see also [5, p. 56]), the following interesting inequality holds:

$$\Phi_{xz} \le \Phi_{xy} + \Phi_{yz}$$
 for any $x, y, z \in H \setminus \{0\}$.

We now recall some inequalities in which the quantity $|\langle x, y \rangle| / (||x|| ||y||)$ for different vectors is involved:

$$(1.4) \qquad 3 \cdot \left| \frac{\langle x, y \rangle \langle y, z \rangle \langle z, x \rangle}{\|x\|^2 \|y\|^2 \|z\|^2} \right| \leq \left| \frac{\langle x, y \rangle}{\|x\| \|y\|} \right|^2 + \left| \frac{\langle y, z \rangle}{\|y\| \|z\|} \right|^2 + \left| \frac{\langle z, x \rangle}{\|z\| \|x\|} \right|^2 \\ \leq 1 + 2 \cdot \left| \frac{\langle x, y \rangle \langle y, z \rangle \langle z, x \rangle}{\|x\|^2 \|y\|^2 \|z\|^2} \right| \qquad [2, p. 37]$$

Date: 19 February, 2008.

²⁰⁰⁰ Mathematics Subject Classification. Primary 47A05, 47A12; Secondary 26D15.

Key words and phrases. Inner product, Schwarz inequality, Bounded linear operators, Integral inequalities, Discrete inequalities.

S. S. DRAGOMIR

(1.5)
$$1 \ge \left| \frac{\langle x, y \rangle}{\|x\| \|y\|} - \frac{\langle x, z \rangle \langle z, y \rangle}{\|x\| \|y\| \|z\|^2} \right| + \left| \frac{\langle x, z \rangle \langle z, y \rangle}{\|x\| \|y\| \|z\|^2} \right|$$
$$\ge \left| \frac{\langle x, y \rangle}{\|x\| \|y\|} \right| \qquad [2, p. 38]$$

$$(1.6) \quad \left| \frac{\langle x, y \rangle \langle y, z \rangle}{\|x\| \|z\| \|y\|^2} \right| \le \frac{1}{2} \cdot \left[1 + \left| \frac{\langle x, z \rangle}{\|x\| \|z\|} \right| \right] \qquad (\text{Buzano's inequality, [2, p. 49]})$$

(1.7)
$$\left|\frac{\langle x, y \rangle \langle y, z \rangle}{\|x\| \|z\| \|y\|^2} - \frac{1}{1+\eta} \cdot \frac{\langle x, z \rangle}{\|x\| \|z\|}\right| \le \frac{1}{\sqrt{2}\sqrt{1+\operatorname{Re}\eta}} \qquad [2, p. 51]$$

where $\eta \in \mathbb{K}$ and $|\eta| = 1$, $\operatorname{Re} \eta \neq -1$, and

$$(1.8) \qquad \left| \frac{\langle x, y \rangle \langle y, z \rangle}{\|x\| \|z\| \|y\|^2} \right| \leq \left| \frac{\langle x, y \rangle \langle y, z \rangle}{\|x\| \|z\| \|y\|^2} - \frac{1}{2} \frac{\langle x, z \rangle}{\|x\| \|z\|} \right| + \frac{1}{2} \cdot \left| \frac{\langle x, z \rangle}{\|x\| \|z\|} \right|$$
$$\leq \frac{1}{2} \cdot \left[1 + \left| \frac{\langle x, z \rangle}{\|x\| \|z\|} \right| \right] \qquad [2, p. 52],$$

where $x, y, z \in H \setminus \{0\}$.

We notice that (1.8) is a refinement of Buzano's inequality (1.6).

For other inequalities of this type, see [1], [4], [7], [8] and [9].

Motivated by the above results, the main aim of the present paper is to compare the quantities

$$\frac{|\langle x,y\rangle|_1}{\|x\|_1 \|y\|_1} \left(\frac{\operatorname{Re}\langle x,y\rangle_1}{\|x\|_1 \|y\|_1}\right) \qquad \text{and} \qquad \frac{|\langle x,y\rangle|_2}{\|x\|_2 \|y\|_2} \left(\frac{\operatorname{Re}\langle x,y\rangle_2}{\|x\|_2 \|y\|_2}\right)$$

in the case when the inner products $\langle \cdot, \cdot \rangle_1$ and $\langle \cdot, \cdot \rangle_2$ defined on H^2 generate two equivalent norms, i.e., we recall that $\|\cdot\|_1$ and $\|\cdot\|_2$ are equivalent if there exists the constants m, M > 0 such that

(1.9)
$$m \|x\|_2 \le \|x\|_1 \le M \|x\|_2$$
, for any $x \in H$.

Applications for invertible bounded linear operators, positive definite self-adjoint operators, integral and discrete inequalities are also given.

2. The Results

The following result may be stated.

Theorem 1. Assume that the inner products $\langle \cdot, \cdot \rangle_i$, $i \in \{1, 2\}$ on the real or complex linear space H generate the norms $\|\cdot\|_i$, $i \in \{1, 2\}$ which satisfy the following condition:

(2.1)
$$m \|x\|_2 \le \|x\|_1 \le M \|x\|_2$$
 for any $x \in H$,

where $0 < m \leq M < \infty$ are given constants.

If $x, y \in H \setminus \{0\}$ satisfy the condition $\operatorname{Re} \langle x, y \rangle_2 \ge 0$, then

(2.2)
$$\frac{m^2}{M^2} - 1 + \frac{\operatorname{Re}\langle x, y \rangle_1}{\|x\|_1 \|y\|_1} \le \frac{\operatorname{Re}\langle x, y \rangle_2}{\|x\|_2 \|y\|_2} \le \frac{\operatorname{Re}\langle x, y \rangle_1}{\|x\|_1 \|y\|_1} + \frac{M^2}{m^2} - 1.$$

If $\operatorname{Re}\langle x,y
angle_2 < 0$, then

(2.3)
$$1 - \frac{m^2}{M^2} + \frac{\operatorname{Re}\langle x, y \rangle_1}{\|x\|_1 \|y\|_1} \le \frac{\operatorname{Re}\langle x, y \rangle_2}{\|x\|_2 \|y\|_2} \le \frac{\operatorname{Re}\langle x, y \rangle_1}{\|x\|_1 \|y\|_1} + 1 - \frac{M^2}{m^2}.$$

Proof. For any inner product $\langle \cdot, \cdot \rangle$ on H, we have

(2.4)
$$1 - \frac{\operatorname{Re}\langle x, y \rangle}{\|x\| \, \|y\|} = \frac{1}{2} \left\| \frac{x}{\|x\|} - \frac{y}{\|y\|} \right\|^2, \quad x, y \in H \setminus \{0\}.$$

Utilising the assumption (2.1), we have successively:

$$(2.5) 1 - \frac{\operatorname{Re}\langle x, y \rangle_1}{\|x\|_1 \|y\|_1} = \frac{1}{2} \left\| \frac{x}{\|x\|_1} - \frac{y}{\|y\|_1} \right\|_1^2 \le \frac{M^2}{2} \left\| \frac{x}{\|x\|_1} - \frac{y}{\|y\|_1} \right\|_2 \\ = \frac{M^2}{2} \left[\frac{\|x\|_2^2}{\|x\|_1^2} + \frac{\|y\|_2^2}{\|y\|_1^2} - \frac{2\operatorname{Re}\langle x, y \rangle_2}{\|x\|_1 \|y\|_1} \right] \\ \le \frac{M^2}{2} \left[\frac{1}{m^2} + \frac{1}{m^2} - 2\frac{\operatorname{Re}\langle x, y \rangle_2}{\|x\|_1 \|y\|_1} \right] \\ = M^2 \left[\frac{1}{m^2} - \frac{\operatorname{Re}\langle x, y \rangle_2}{\|x\|_1 \|y\|_1} \right] =: I.$$

Now, if $\operatorname{Re}\langle x,y\rangle_2 \ge 0$, then

$$-\frac{\operatorname{Re}\langle x, y \rangle_2}{\|x\|_1 \|y\|_1} \le -\frac{\operatorname{Re}\langle x, y \rangle_2}{M^2 \|x\|_2 \|y\|_2}$$

which implies that

(2.6)
$$I \le M^2 \left[\frac{1}{m^2} - \frac{\operatorname{Re} \langle x, y \rangle_2}{M^2 \|x\|_2 \|y\|_2} \right] = \frac{M^2}{m^2} - \frac{\operatorname{Re} \langle x, y \rangle_2}{\|x\|_2 \|y\|_2}.$$

Utilising (2.5) and (2.6) we deduce

$$1 - \frac{\operatorname{Re} \langle x, y \rangle_1}{\|x\|_1 \|y\|_1} \le \frac{M^2}{m^2} - \frac{\operatorname{Re} \langle x, y \rangle_2}{\|x\|_2 \|y\|_2},$$

which produces the second inequality in (2.2).

By (2.4) and (2.1) we also have

$$(2.7) 1 - \frac{\operatorname{Re}\langle x, y \rangle_1}{\|x\|_1 \|y\|_1} \ge \frac{m^2}{2} \left\| \frac{x}{\|x\|_1} - \frac{y}{\|y\|_1} \right\|_2^2 \\ = \frac{m^2}{2} \left[\frac{\|x\|_2^2}{\|x\|_1^2} + \frac{\|y\|_2^2}{\|y\|_1^2} - 2\frac{\operatorname{Re}\langle x, y \rangle_2}{\|x\|_1 \|y\|_1} \right] \\ \ge \frac{m^2}{2} \left[\frac{1}{M^2} + \frac{1}{M^2} - 2\frac{\operatorname{Re}\langle x, y \rangle_2}{\|x\|_1 \|y\|_1} \right] \\ = m^2 \left[\frac{1}{M^2} - \frac{\operatorname{Re}\langle x, y \rangle_2}{\|x\|_1 \|y\|_1} \right] =: J.$$

Due to the fact that $\operatorname{Re}\langle x,y\rangle_2 \geq 0$, we have

$$-\frac{\operatorname{Re} \langle x, y \rangle_2}{\|x\|_1 \|y\|_1} \ge -\frac{\operatorname{Re} \langle x, y \rangle_2}{m^2 \|x\|_2 \|y\|_2},$$

which implies that

(2.8)
$$J \ge m^2 \left[\frac{1}{M^2} - \frac{\operatorname{Re} \langle x, y \rangle_2}{m^2 \|x\|_2 \|y\|_2} \right] = \frac{m^2}{M^2} - \frac{\operatorname{Re} \langle x, y \rangle_2}{\|x\|_2 \|y\|_2}$$

By making use of (2.7) and (2.8) we get

$$1 - \frac{\operatorname{Re} \langle x, y \rangle_1}{\|x\|_1 \|y\|_1} \ge \frac{m^2}{M^2} - \frac{\operatorname{Re} \langle x, y \rangle_2}{\|x\|_2 \|y\|_2},$$

which is clearly equivalent to the first inequality in (2.2).

Finally, if $\operatorname{Re} \langle x, y \rangle_2 < 0$, then $\operatorname{Re} \langle x, -y \rangle_2 > 0$ and writing the inequality (2.2) for -y instead of y, we easily deduce (2.3).

Corollary 1. Let $A \in B(H)$ be an invertible operator on the Hilbert space $(H; \langle \cdot, \cdot \rangle)$. If $x, y \in H \setminus \{0\}$ are such that $\operatorname{Re} \langle x, y \rangle \geq 0$, then:

(2.9)
$$1 - \|A\|^{2} \|A^{-1}\|^{2} + \frac{\operatorname{Re}\langle x, y \rangle}{\|x\| \|y\|} \leq \frac{\operatorname{Re}\langle Ax, Ay \rangle}{\|Ax\| \|Ay\|} \leq \frac{\operatorname{Re}\langle x, y \rangle}{\|x\| \|y\|} + 1 - \frac{1}{\|A\|^{2} \|A^{-1}\|^{2}}$$

If $x, y \in H \setminus \{0\}$ and $\operatorname{Re} \langle x, y \rangle < 0$, then

(2.10)
$$\|A\|^{2} \|A^{-1}\|^{2} - 1 + \frac{\operatorname{Re}\langle x, y \rangle}{\|x\| \|y\|} \leq \frac{\operatorname{Re}\langle Ax, Ay \rangle}{\|Ax\| \|Ay\|} \\ \leq \frac{\operatorname{Re}\langle x, y \rangle}{\|x\| \|y\|} + \frac{1}{\|A\|^{2} \|A^{-1}\|^{2}} - 1.$$

Proof. Since $A \in B(H)$ is invertible, then

$$\frac{1}{\|A^{-1}\|} \cdot \|x\| \le \|Ax\| \le \|A\| \|x\|, \quad \text{for any } x \in H.$$

Applying Theorem 1 for $\langle x, y \rangle_1 := \langle Ax, Ay \rangle$, $\langle x, y \rangle_2 := \langle x, y \rangle$ and $m = \frac{1}{\|A^{-1}\|}$, $M = \|A\|$ and doing the necessary calculations, we deduce the desired result. \Box

Corollary 2. Let $A \in B(H)$ be a self-adjoint operator on the Hilbert space $(H; \langle \cdot, \cdot \rangle)$ which satisfies the condition

(2.11)
$$\gamma I \le A \le \Gamma I,$$

in the operator order of B(H), where $0 < \gamma \leq \Gamma < \infty$ are given. If $x, y \in H \setminus \{0\}$ are such that $\operatorname{Re} \langle x, y \rangle \geq 0$, then

(2.12)
$$1 - \frac{\Gamma}{\gamma} + \frac{\operatorname{Re}\langle x, y \rangle}{\|x\| \, \|y\|} \le \frac{\operatorname{Re}\langle Ax, y \rangle}{\left[\langle Ax, x \rangle \, \langle Ay, y \rangle\right]^{\frac{1}{2}}} \le \frac{\operatorname{Re}\langle x, y \rangle}{\|x\| \, \|y\|} + 1 - \frac{\gamma}{\Gamma}.$$

If $x, y \in H \setminus \{0\}$ are such that $\operatorname{Re} \langle x, y \rangle < 0$, then

(2.13)
$$\frac{\Gamma}{\gamma} - 1 + \frac{\operatorname{Re}\langle x, y \rangle}{\|x\| \|y\|} \le \frac{\operatorname{Re}\langle Ax, y \rangle}{[\langle Ax, x \rangle \langle Ay, y \rangle]^{\frac{1}{2}}} \le \frac{\operatorname{Re}\langle x, y \rangle}{\|x\| \|y\|} + \frac{\gamma}{\Gamma} - 1.$$

Proof. From (2.11) we have

 $\gamma \langle x, x \rangle \le \langle Ax, x \rangle \le \Gamma \langle x, x \rangle, \qquad x \in H,$

which implies that $\sqrt{\gamma} \|x\| \le [\langle Ax, x \rangle]^{1/2} \le \sqrt{\Gamma} \|x\|, x \in H.$

Now, if we apply Theorem 1 for $\langle x, y \rangle_1 := \langle Ax, y \rangle$, $\langle x, y \rangle_2 := \langle x, y \rangle$, $x, y \in H$ and $m = \sqrt{\gamma}$, $M = \sqrt{\Gamma}$, then we obtain the desired result.

The following lemma is of interest in itself.

Lemma 1. Assume that the inner products $\langle \cdot, \cdot \rangle_i$, $i \in \{1, 2\}$ defined on H satisfy the condition (2.1). Then for any $x, y \in H$ we have

(2.14)
$$m^{2} [\|x\|_{2} \|y\|_{2} - |\langle x, y \rangle|_{2}] \leq \|x\|_{1} \|y\|_{1} - |\langle x, y \rangle|_{1} \\ \leq M^{2} [\|x\|_{2} \|y\|_{2} - |\langle x, y \rangle|_{2}]$$

and

(2.15)
$$m^{2} [\|x\|_{2} \|y\|_{2} - \operatorname{Re} \langle x, y \rangle_{2}] \leq \|x\|_{1} \|y\|_{1} - \operatorname{Re} \langle x, y \rangle_{1} \\ \leq M^{2} [\|x\|_{2} \|y\|_{2} - \operatorname{Re} \langle x, y \rangle_{2}],$$

respectively.

Proof. We use the following result obtained by Dragomir and Mond in [3] (see also [2, p. 9]):

If $[\cdot, \cdot]_1$, $[\cdot, \cdot]_2$ are two hermitian forms on H with $[x, x]_1^{1/2} \leq [x, x]_2^{1/2}$ for any $x \in H$, then

(2.16)
$$[x,x]_1^{1/2} [y,y]_1^{1/2} - |[x,y]_1| \le [x,x]_2^{1/2} [y,y]_2^{1/2} - |[x,y]_2|$$

and

(2.17)
$$[x, x]_{1}^{1/2} [y, y]_{1}^{1/2} - \operatorname{Re}[x, y]_{1} \le [x, x]_{2}^{1/2} [y, y]_{2}^{1/2} - \operatorname{Re}[x, y]_{2}$$

for any $x, y \in H$.

Now, if we apply (2.16) and (2.17) firstly for $[\cdot, \cdot]_2 := \langle \cdot, \cdot \rangle_1$, $[\cdot, \cdot]_1 = m^2 \langle \cdot, \cdot \rangle_2$ and then for $[\cdot, \cdot]_2 = M^2 \langle \cdot, \cdot \rangle_2$, $[\cdot, \cdot]_1 := \langle \cdot, \cdot \rangle_1$, we deduce the desired results (2.14) and (2.15).

The following result may be stated as well.

Theorem 2. Assume that the inner products $\langle \cdot, \cdot \rangle_i$, $i \in \{1, 2\}$ satisfy the condition (2.1). Then for any $x, y \in H \setminus \{0\}$, we have the inequalities:

$$(2.18) \qquad \frac{m^2}{M^2} - 1 + \frac{|\langle x, y \rangle_1|}{\|x\|_1 \|y\|_1} \le \frac{|\langle x, y \rangle_2|}{\|x\|_2 \|y\|_2} \le \frac{|\langle x, y \rangle_1|}{\|x\|_1 \|y\|_1} + \frac{M^2}{m^2} - 1$$

Proof. Dividing the inequality (2.14) by $||x||_1 ||y||_1 \neq 0$, we obtain

$$(2.19) mtext{ } m^{2} \left[\frac{\|x\|_{2} \|y\|_{2}}{\|x\|_{1} \|y\|_{1}} - \frac{|\langle x, y \rangle_{2}|}{\|x\|_{1} \|y\|_{1}} \right] \leq 1 - \frac{|\langle x, y \rangle_{1}|}{\|x\|_{1} \|y\|_{1}} \\ \leq M^{2} \left[\frac{\|x\|_{2} \|y\|_{2}}{\|x\|_{1} \|y\|_{1}} - \frac{|\langle x, y \rangle_{2}|}{\|x\|_{1} \|y\|_{1}} \right],$$

for any $x, y \in H \setminus \{0\}$.

Observe, by (2.1) that:

$$\frac{\|x\|_2 \, \|y\|_2}{\|x\|_1 \, \|y\|_1} \leq \frac{1}{m^2} \quad \text{and} \quad - \frac{|\langle x, y \rangle_2|}{\|x\|_1 \, \|y\|_1} \leq -\frac{|\langle x, y \rangle_2|}{M^2 \, \|x\|_2 \, \|y\|_2}$$

Utilising the second inequality in (2.19), we deduce

$$1 - \frac{|\langle x, y \rangle_1|}{\|x\|_1 \|y\|_1} \le \frac{M^2}{m^2} - \frac{|\langle x, y \rangle_2|}{\|x\|_2 \|y\|_2},$$

which is equivalent with the second inequality in (2.18).

In addition, we have

$$\frac{1}{M^2} \leq \frac{\|x\|_2 \|y\|_2}{\|x\|_1 \|y\|_1} \quad \text{and} \quad -\frac{|\langle x, y \rangle_2|}{m^2 \|x\|_2 \|y\|_2} \leq -\frac{|\langle x, y \rangle_1|}{\|x\|_1 \|y\|_1}$$

which together with the first inequality in (2.19) produces the first part of (2.18). \Box

Remark 1. On utilising the inequality (2.15) and an argument similar to the one in the proof of Theorem 2, we can reobtain the inequalities (2.2) and (2.3) from Theorem 1. The details are omitted.

Corollary 3. Let $A \in B(H)$ be an invertible operator on the Hilbert space $(H; \langle \cdot, \cdot \rangle)$. Then

(2.20)
$$1 - \|A\|^{2} \|A^{-1}\|^{2} + \frac{|\langle x, y \rangle|}{\|x\| \|y\|} \leq \frac{|\langle Ax, Ay \rangle|}{\|Ax\| \|Ay\|} \leq \frac{|\langle x, y \rangle|}{\|x\| \|y\|} + 1 - \frac{1}{\|A\|^{2} \|A^{-1}\|^{2}}$$

for any $x, y \in H \setminus \{0\}$.

The proof follows from Theorem 2 on choosing $\langle x, y \rangle_1 := \langle Ax, Ay \rangle$, $\langle x, y \rangle_2 := \langle x, y \rangle$, $x, y \in H$ and $m = \frac{1}{\|A^{-1}\|}$, $M = \|A\|$.

Corollary 4. Let $A \in B(H)$ be a self-adjoint operator satisfying the condition (2.11). Then

$$(2.21) 1 - \frac{\Gamma}{\gamma} + \frac{|\langle x, y \rangle|}{\|x\| \|y\|} \le \frac{|\langle Ax, y \rangle|}{[\langle Ax, x \rangle \langle Ay, y \rangle]^{\frac{1}{2}}} \le \frac{|\langle x, y \rangle|}{\|x\| \|y\|} + 1 - \frac{\gamma}{\Gamma},$$

for any $x, y \in H \setminus \{0\}$.

The proof follows by Theorem 2 on choosing $\langle x, y \rangle_1 := \langle Ax, y \rangle$, $\langle x, y \rangle_2 := \langle x, y \rangle$, $x, y \in H$ and $m = \sqrt{\gamma}$, $M = \sqrt{\Gamma}$.

3. Applications for Integral Inequalities

Assume that $(K; \langle \cdot, \cdot \rangle)$ is a Hilbert space over the real or complex number field \mathbb{K} . If $\rho : [a,b] \subset \mathbb{R} \to [0,\infty)$ is a measurable function then we may consider the space $L^2_{\rho}([a,b];K)$ of all functions $f:[a,b] \to K$ that are strongly measurable and $\int_a^b \rho(t) \|f(t)\|^2 dt < \infty$. It is well known that $L^2_{\rho}([a,b];K)$ endowed with the inner product $\langle \cdot, \cdot \rangle_{\rho}$ defined by

$$\left\langle f,g
ight
angle _{
ho}:=\int_{a}^{b}
ho\left(t
ight) \left\langle f\left(t
ight) ,g\left(t
ight)
ight
angle dt$$

and generating the norm

$$\|f\|_{\rho} := \left(\int_{a}^{b} \rho\left(t\right) \|f\left(t\right)\|^{2} dt\right)^{\frac{1}{2}}$$

is a Hilbert space over $\mathbbm{K}.$

The following proposition can be stated.

Proposition 1. Let $\rho_1, \rho_2 : [a, b] \to [0, \infty)$ be two measurable functions with the property that there exists $0 < \phi \leq \Phi < \infty$ so that

(3.1)
$$\phi \leq \frac{\rho_1(t)}{\rho_2(t)} \leq \Phi \quad for \ a.e. \ t \in [a,b]$$

Then for any $f, g \in L^2_{\rho_2}([a, b]; K)$ we have the inequalities:

$$(3.2) \qquad \frac{\phi}{\Phi} - 1 + \frac{\left|\int_{a}^{b} \rho_{1}\left(t\right) \left\langle f\left(t\right), g\left(t\right) \right\rangle dt\right|}{\left(\int_{a}^{b} \rho_{1}\left(t\right) \left\|f\left(t\right)\right\|^{2} dt\right)^{\frac{1}{2}} \left(\int_{a}^{b} \rho_{2}\left(t\right) \left\|g\left(t\right)\right\|^{2} dt\right)^{\frac{1}{2}}} \\ \leq \frac{\left|\int_{a}^{b} \rho_{2}\left(t\right) \left\langle f\left(t\right), g\left(t\right) \right\rangle dt\right|}{\left(\int_{a}^{b} \rho_{2}\left(t\right) \left\|f\left(t\right)\right\|^{2} dt\right)^{\frac{1}{2}} \left(\int_{a}^{b} \rho_{2}\left(t\right) \left\|g\left(t\right)\right\|^{2} dt\right)^{\frac{1}{2}}} \\ \leq \frac{\left|\int_{a}^{b} \rho_{1}\left(t\right) \left\langle f\left(t\right), g\left(t\right) \right\rangle dt\right|}{\left(\int_{a}^{b} \rho_{1}\left(t\right) \left\|f\left(t\right)\right\|^{2} dt\right)^{\frac{1}{2}} \left(\int_{a}^{b} \rho_{2}\left(t\right) \left\|g\left(t\right)\right\|^{2} dt\right)^{\frac{1}{2}}} + \frac{\Phi}{\phi} - 1$$

Proof. From (3.1) we have

$$\begin{split} \sqrt{\phi} \left(\int_{a}^{b} \rho_{2}\left(t\right) \left\|f\left(t\right)\right\|^{2} dt \right)^{\frac{1}{2}} &\leq \left(\int_{a}^{b} \rho_{1}\left(t\right) \left\|f\left(t\right)\right\|^{2} dt \right)^{\frac{1}{2}} \\ &\leq \sqrt{\Phi} \left(\int_{a}^{b} \rho_{2}\left(t\right) \left\|f\left(t\right)\right\|^{2} dt \right)^{\frac{1}{2}}. \end{split}$$

Applying Theorem 2 for $\langle \cdot, \cdot \rangle_i = \langle \cdot, \cdot \rangle_{\rho_i}$, $i \in \{1, 2\}$ and $H = L^2_{\rho_1}([a, b]; K) = L^2_{\rho_2}([a, b]; K)$, we deduce the desired result.

Remark 2. A similar result can be stated if one uses Theorem 1. The details are omitted.

4. Applications for Discrete Inequalities

Assume that $(K; \langle \cdot, \cdot \rangle)$ is a Hilbert space over the real or complex number field and $\mathbf{p} = (p_i)_{i \in \mathbb{N}}$ with $p_i \ge 0$, $i \in \mathbb{N}$ and $\sum_{i=1}^{\infty} p_i < \infty$. Define

$$\ell_p^2(K) := \left\{ \mathbf{x} := (x_i)_{i \in \mathbb{N}} \mid x_i \in K, \ i \in \mathbb{N} \text{ and } \sum_{i=1}^{\infty} p_i \|x_i\|^2 < \infty \right\}.$$

It is well known that $\ell_p^2(K)$ endowed with the inner product

$$\langle \mathbf{x}, \mathbf{y} \rangle_p := \sum_{i=1}^n p_i \langle x_i, y_i \rangle$$

and generating the norm

$$\|\mathbf{x}\|_p := \left(\sum_{i=1}^{\infty} p_i \|x_i\|^2\right)^{\frac{1}{2}}$$

is a Hilbert space over \mathbb{K} .

Proposition 2. Assume that $\mathbf{p} = (p_i)_{i \in \mathbb{N}}$, $\mathbf{q} = (q_i)_{i \in \mathbb{N}}$ are such that $p_i, q_i \ge 0$, $i \in \mathbb{N}$, $\sum_{i=1}^{\infty} p_i, \sum_{i=1}^{\infty} q_i < \infty$ and

$$(4.1) nq_i \le p_i \le Nq_i for any i \in \mathbb{N},$$

where $0 < n \leq N < \infty$. Then we have the inequality

(4.2)
$$\frac{n}{N} - 1 + \frac{\left|\sum_{i=1}^{\infty} p_i \langle x_i, y_i \rangle\right|}{\left(\sum_{i=1}^{\infty} p_i \|x_i\|^2\right)^{\frac{1}{2}} \left(\sum_{i=1}^{\infty} p_i \|y_i\|^2\right)^{\frac{1}{2}}} \\ \leq \frac{\left|\sum_{i=1}^{\infty} q_i \langle x_i, y_i \rangle\right|}{\left(\sum_{i=1}^{\infty} q_i \|x_i\|^2\right)^{\frac{1}{2}} \left(\sum_{i=1}^{\infty} q_i \|y_i\|^2\right)^{\frac{1}{2}}} \\ \leq \frac{\left|\sum_{i=1}^{\infty} p_i \langle x_i, y_i \rangle\right|}{\left(\sum_{i=1}^{\infty} p_i \|x_i\|^2\right)^{\frac{1}{2}} \left(\sum_{i=1}^{\infty} p_i \|y_i\|^2\right)^{\frac{1}{2}}} + \frac{N}{n} - 1.$$

A similar result can be stated if one uses Theorem 1. However the details are omitted.

References

- M.L. BUZANO, Generalizazione della diseguaglianza di Cauchy-Schwarz (Italian), Rend. Sem. Mat. Univ. e Politech. Torino, 31 (1971/73), 405-409.
- [2] S.S. DRAGOMIR, Advances in Inequalities of the Schwarz, Triangle and Heisenberg Type in Inner Product Spaces, Nova Science Publishers, NY, 2007.
- [3] S.S. DRAGOMIR and B. MOND, On the superadditivity and monotonicity of Schwarz's inequality in inner product spaces, *Makedon. Akad. Nauk. Umet. Oddel. Mat.-Tehn. Nauk. Prilozi* 15 (1994), no. 2, 5–22 (1996).
- [4] S.S. DRAGOMIR and J. SÁNDOR, Some inequalities in prehilbertian spaces, Studia Univ. Babeş-Bolyai, Math., 32(11) (1987), 71-78.
- [5] K.E. GUSTAFSON and D.K.M. RAO, Numerical Range, Springer, 1997.
- [6] M. KREIN, Angular localization of the spectrum of a multiplicative integral in a Hilbert space, Funct. Anal. Appl., 3 (1969), 89-90.
- [7] M.H. MOORE, An inner product inequality, SIAM J. Math. Anal., 4(3) (1973), 514-518.
- [8] T. PRECUPANU, On a generalization of Cauchy-Buniakowski-Schwarz inequality, Anal. St. Univ. "Al. I. Cuza" Iaşi, 22(2) (1976), 173-175.
- [9] U. RICHARD, Sur des inégalités du type Wirtinger et leurs application aux équationes différentiable ordinaires, Coll. of Anal. Rio de Janeiro, August, 1972, 233-244.

School of Computer Science and Mathematics, Victoria University, PO Box 14428, Mebourne City 8001, VICTORIA, Australia.

E-mail address: sever.dragomir@vu.edu.au *URL*: http://rgmia.vu.edu.au/dragomir