



**VICTORIA UNIVERSITY**  
MELBOURNE AUSTRALIA

## *Jensen's and Hermite–Hadamard's Type Inequalities for Lower and Strongly Convex Functions on Normed Spaces*

This is the Published version of the following publication

Dragomir, Sever S and Nikodem, K (2018) Jensen's and Hermite–Hadamard's Type Inequalities for Lower and Strongly Convex Functions on Normed Spaces. *Bulletin of the Iranian Mathematical Society*, 44 (5). 1337 - 1349. ISSN 1017-060X

The publisher's official version can be found at  
<https://link.springer.com/article/10.1007/s41980-018-0095-9>  
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/37809/>



# Jensen's and Hermite–Hadamard's Type Inequalities for Lower and Strongly Convex Functions on Normed Spaces

Silvestru Sever Dragomir<sup>1,2</sup> · Kazimierz Nikodem<sup>3</sup>

Received: 4 June 2017 / Accepted: 25 December 2017 / Published online: 21 June 2018

© The Author(s) 2018

## Abstract

In this paper, we obtain some Jensen's and Hermite–Hadamard's type inequalities for lower, upper, and strongly convex functions defined on convex subsets in normed linear spaces. The case of inner product space is of interest since in these case the concepts of lower convexity and strong convexity coincides. Applications for univariate functions of real variable and the connections with earlier Hermite–Hadamard's type inequalities are also provided.

**Keywords** Jensen's inequality · Hermite–Hadamard inequality · Strongly convex functions · Lower convex functions

**Mathematics Subject Classification** Primary 26A51; Secondary 26D15 · 39B62

## 1 Introduction

The classical Jensen inequality and Hermite–Hadamard inequality are ones of the most important inequalities in convex analysis and they have various applications in mathematics, statistics, economics and engineering sciences. The aim of this paper is to present counterparts of these inequalities for  $m$ -lower convex and strongly convex

---

Communicated by Ali Ghaffari.

---

✉ Kazimierz Nikodem  
knikodem@ath.bielsko.pl  
Silvestru Sever Dragomir  
sever.dragomir@vu.edu.au

<sup>1</sup> Mathematics, College of Engineering and Science, Victoria University, P.O. Box 14428, Melbourne VIC, 8001, Australia

<sup>2</sup> DST-NRF Centre of Excellence in the Mathematical and Statistical Sciences, School of Computer Science and Applied Mathematics, University of the Witwatersrand, Wits, Private Bag 3, Johannesburg 2050, South Africa

<sup>3</sup> Department of Mathematics, University of Bielsko-Biala, ul. Willowa 2, 43-309 Bielsko-Biala, Poland

functions in normed spaces. The theorem obtained by us extend and complement some earlier results on Jensen and Hermite–Hadamard inequalities obtained, among others, in [3,7,13,14,16,19].

Let  $X$  be a real linear space. Assume that  $a, b \in X$ ,  $a \neq b$  and let  $[a, b] := \{(1 - \lambda)a + \lambda b, \lambda \in [0, 1]\}$  be the *segment* generated by  $a$  and  $b$ . We consider the function  $f : [a, b] \rightarrow \mathbb{R}$  and the attached function  $g(a, b) : [0, 1] \rightarrow \mathbb{R}$ ,  $g(a, b)(t) := f[(1 - t)a + tb]$ ,  $t \in [0, 1]$ .

It is well known that  $f$  is convex on  $[a, b]$  iff  $g(a, b)$  is convex on  $[0, 1]$ , and the following lateral derivatives exist and satisfy the properties:

- (i)  $g'_{\pm}(a, b)(s) = (\nabla_{\pm} f[(1 - s)a + sb])(b - a)$ ,  $s \in (0, 1)$ ;
- (ii)  $g'_{+}(a, b)(0) = (\nabla_{+} f(a))(b - a)$ ;
- (iii)  $g'_{-}(a, b)(1) = (\nabla_{-} f(b))(b - a)$ ;

where  $(\nabla_{\pm} f(x))(y)$  are the *Gâteaux lateral derivatives*. Recall that

$$(\nabla_{+} f(x))(y) := \lim_{h \rightarrow 0+} \left[ \frac{f(x + hy) - f(x)}{h} \right],$$

$$(\nabla_{-} f(x))(y) := \lim_{k \rightarrow 0-} \left[ \frac{f(x + ky) - f(x)}{k} \right], \quad x, y \in X.$$

Now, assume that  $(X, \|\cdot\|)$  is a normed linear space. The function  $f_0(x) = \frac{1}{2} \|x\|^2$ ,  $x \in X$  is convex, and thus, the following limits exist

- (iv)  $\langle x, y \rangle_s := (\nabla_{+} f_0(y))(x) = \lim_{t \rightarrow 0+} \left[ \frac{\|y + tx\|^2 - \|y\|^2}{2t} \right]$ ;
- (v)  $\langle x, y \rangle_i := (\nabla_{-} f_0(y))(x) = \lim_{s \rightarrow 0-} \left[ \frac{\|y + sx\|^2 - \|y\|^2}{2s} \right]$ ;

for any  $x, y \in X$ . They are called the *lower* and *upper semi-inner products* associated to the norm  $\|\cdot\|$ , respectively.

For the sake of completeness, we list here some of the main properties of these mappings that will be used in the sequel (see for example [6]), assuming that  $p, q \in \{s, i\}$  and  $p \neq q$ . Recall also that a function  $f : X \rightarrow \mathbb{R}$  is called *subadditive* (*superadditive*) if  $f(x_1 + x_2) \leq f(x_1) + f(x_2)$  ( $f(x_1 + x_2) \geq f(x_1) + f(x_2)$ ) for all  $x_1, x_2 \in X$ .

- (a)  $\langle x, x \rangle_p = \|x\|^2$  for all  $x \in X$ ;
- (aa)  $\langle \alpha x, \beta y \rangle_p = \alpha \beta \langle x, y \rangle_p$  if  $\alpha, \beta \geq 0$  and  $x, y \in X$ ;
- (aaa)  $|\langle x, y \rangle_p| \leq \|x\| \|y\|$  for all  $x, y \in X$ ;
- (av)  $\langle \alpha x + y, x \rangle_p = \alpha \langle x, x \rangle_p + \langle y, x \rangle_p$  if  $x, y \in X$  and  $\alpha \in \mathbb{R}$ ;
- (v)  $\langle -x, y \rangle_p = -\langle x, y \rangle_q$  for all  $x, y \in X$ ;
- (va)  $\langle x + y, z \rangle_p \leq \|x\| \|z\| + \langle y, z \rangle_p$  for all  $x, y, z \in X$ ;
- (vaa) The mapping  $\langle \cdot, \cdot \rangle_p$  is continuous and subadditive (superadditive) in the first variable for  $p = s$  (or  $p = i$ );
- (vaaa) If the norm  $\|\cdot\|$  is induced by an inner product  $\langle \cdot, \cdot \rangle$ , then  $\langle y, x \rangle_i = \langle y, x \rangle = \langle y, x \rangle_s$  for all  $x, y \in X$ .

The following inequality is the well-known Hermite–Hadamard integral inequality for convex functions defined on a segment  $[a, b] \subset X$ :

$$f\left(\frac{a+b}{2}\right) \leq \int_0^1 f[(1-t)a + tb] dt \leq \frac{f(a) + f(b)}{2}, \quad (\text{HH})$$

which easily follows by the classical Hermite–Hadamard inequality for the convex function  $g(a, b) : [0, 1] \rightarrow \mathbb{R}$ . For other related results, see the monograph [14].

Applying inequality (HH) for the convex function  $f_0(x) = \|x\|^2$ , one may deduce the inequality

$$\left\|\frac{x+y}{2}\right\|^2 \leq \int_0^1 \|(1-t)x + ty\|^2 dt \leq \frac{\|x\|^2 + \|y\|^2}{2} \quad (1.1)$$

for any  $x, y \in X$ . The same (HH) inequality applied for  $f_1(x) = \|x\|$ , will give the following refinement of the triangle inequality:

$$\left\|\frac{x+y}{2}\right\| \leq \int_0^1 \|(1-t)x + ty\| dt \leq \frac{\|x\| + \|y\|}{2}, \quad x, y \in X.$$

The distance between the first and second term in (1.1) has the lower and upper bounds [9]:

$$\begin{aligned} 0 &\leq \frac{1}{8} [\langle y-x, y+x \rangle_s - \langle y-x, y+x \rangle_i] \\ &\leq \int_0^1 \|(1-t)x + ty\|^2 dt - \left\|\frac{x+y}{2}\right\|^2 \leq \frac{1}{4} [\langle y-x, y \rangle_i - \langle y-x, x \rangle_s] \end{aligned} \quad (1.2)$$

while the distance between the second and third term in (1.1) has the same upper and lower bounds, namely [10]

$$\begin{aligned} 0 &\leq \frac{1}{8} [\langle y-x, y+x \rangle_s - \langle y-x, y+x \rangle_i] \\ &\leq \frac{\|x\|^2 + \|y\|^2}{2} - \int_0^1 \|(1-t)x + ty\|^2 dt \leq \frac{1}{4} [\langle y-x, y \rangle_i - \langle y-x, x \rangle_s] \end{aligned} \quad (1.3)$$

for any  $x, y \in X$ . The multiplicative constants  $\frac{1}{8}$  and  $\frac{1}{4}$  are best possible in (1.2) and (1.3).

## 2 Some Jensen's Type Inequalities

Let  $(X, \|\cdot\|)$  be a real or complex normed linear space,  $C \subseteq X$  a convex subset of  $X$  and  $f : C \rightarrow \mathbb{R}$ . Let  $m, M \in \mathbb{R}$ . The mapping  $f$  will be called *m-lower convex* on

$C$  if  $f - \frac{m}{2} \|\cdot\|^2$  is a convex mapping on  $C$ . The mapping  $f$  will be called *M-upper convex* on  $C$  if  $\frac{M}{2} \|\cdot\|^2 - f$  is a convex mapping on  $C$ . The mapping  $f$  will be called *(m, M)-convex* on  $C$  if it is both *m-lower convex* and *M-upper convex* on  $C$ . Note that if  $f$  is *(m, M)-convex* on  $C$ , then  $m \leq M$ .

Furthermore, assume that  $c$  is a positive constant. A function  $f : C \rightarrow \mathbb{R}$  is called: *strongly convex with modulus c* if

$$f(tx + (1-t)y) \leq tf(x) + (1-t)f(y) - ct(1-t)\|x - y\|^2 \quad (2.1)$$

for all  $x, y \in C$  and  $t \in [0, 1]$ . In addition, it is called: *strongly Jensen-convex with modulus c* if (2.1) is assumed only for  $t = \frac{1}{2}$ , that is

$$f\left(\frac{x+y}{2}\right) \leq \frac{f(x) + f(y)}{2} - \frac{c}{4}\|x - y\|^2, \quad \text{for all } x, y \in C.$$

The usual concepts of convexity and Jensen-convexity correspond to the case  $c = 0$ , respectively. The notion of strongly convex functions have been introduced by Polyak [23] and they play an important role in optimization theory and mathematical economics. Many properties and applications of them can be found in the literature (see, for instance, [1,2,5,15–18,22–26]). Let us mention also the paper [21] by the second author which is a survey article devoted to strongly convex functions and related classes of functions.

Denote by  $\mathcal{SC}(C, c)$  the class of all functions  $f : C \rightarrow \mathbb{R}$  strongly convex with modulus  $c$  and by  $\mathcal{LC}(C, m)$  the class of all functions  $f : C \rightarrow \mathbb{R}$  *m-lower convex*. It is known that [22] if  $X$  is an inner product space then

$$\mathcal{SC}\left(C, \frac{m}{2}\right) = \mathcal{LC}(C, m).$$

However, in arbitrary normed spaces the above classes differ in general.

The following examples shows that neither  $\mathcal{LC}(C, m)$  is included in  $\mathcal{SC}\left(C, \frac{m}{2}\right)$ , nor conversely.

**Example 2.1** ([22]) Let  $X = \mathbb{R}^2$  and  $\|x\| = |x_1| + |x_2|$ , for  $x = (x_1, x_2)$ . Take  $f = \|\cdot\|^2$ . Then  $g = f - \|\cdot\|^2$  is convex being the zero function. However,  $f$  is not strongly convex with modulus 1. Indeed, for  $x = (1, 0)$  and  $y = (0, 1)$ , we have

$$f\left(\frac{x+y}{2}\right) = 1 > 0 = \frac{f(x) + f(y)}{2} - \frac{1}{4}\|x - y\|^2,$$

which contradicts (2.1).

**Example 2.2** Let  $X = \mathbb{R}^2$  and  $\|x\| = |x_1| + |x_2|$ , for  $x = (x_1, x_2)$ . Take  $f(x) = x_1^2 + x_2^2$ . Then,  $f$  is strongly convex with modulus  $c = \frac{1}{2}$ . Indeed, for arbitrary  $x = (x_1, x_2)$  and  $y = (y_1, y_2)$  in  $\mathbb{R}^2$ , we have

$$f\left(\frac{x+y}{2}\right) = \frac{1}{4}\left(x_1^2 + 2x_1y_1 + y_1^2 + x_2^2 + 2x_2y_2 + y_2^2\right)$$

and

$$\begin{aligned} & \frac{f(x) + f(y)}{2} - \frac{1}{2} \cdot \frac{1}{4} \|x - y\|^2 \\ &= \frac{3}{8} (x_1^2 + y_1^2 + x_2^2 + y_2^2) + \frac{1}{4} (x_1 y_1 + x_2 y_2 - |x_1 - y_1| |x_2 - y_2|). \end{aligned}$$

Hence

$$\frac{f(x) + f(y)}{2} - \frac{1}{2} \cdot \frac{1}{4} \|x - y\|^2 - f\left(\frac{x + y}{2}\right) = \frac{1}{8} (|x_1 - y_1| - |x_2 - y_2|)^2 \geq 0.$$

This shows that  $f$  is strongly Jensen-convex with modulus  $c = \frac{1}{2}$ . Since  $f$  is continuous, it is also strongly convex with modulus  $c = \frac{1}{2}$ . On the other hand, the function  $g = f - \frac{1}{2} \|\cdot\|^2$  is not convex. Indeed, for  $x = (-1, 1)$  and  $y = (1, 1)$ , we have

$$g\left(\frac{x + y}{2}\right) = \frac{1}{2} > 0 = \frac{g(x) + g(y)}{2}.$$

The following Jensen's type inequality holds [7].

**Proposition 2.3** *Let  $f : C \subseteq X \rightarrow \mathbb{R}$ ,  $C$  be convex on  $X$ ,  $x_i \in C$ ,  $p_i \geq 0$  ( $i = 1, \dots, n$ ) with  $\sum_{i=1}^n p_i = 1$ . If  $f$  is  $m$ -lower convex on  $C$ , then we have the following inequality (for  $m \geq 0$  - refinement of Jensen's inequality)*

$$\frac{m}{2} \left[ \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 \right] \leq \sum_{i=1}^n p_i f(x_i) - f\left(\sum_{i=1}^n p_i x_i\right). \quad (2.2)$$

As a consequence of the above proposition, we get the following corollary for functions defining on inner product spaces.

**Corollary 2.4** *Let  $(X, \langle \cdot, \cdot \rangle)$  be an inner product space,  $\|\cdot\| := \sqrt{\langle \cdot, \cdot \rangle}$ ,  $C \subseteq X$  a convex subset on  $X$ ,  $f : C \rightarrow \mathbb{R}$  and  $x_i \in C$ ,  $p_i \geq 0$  ( $i = 1, \dots, n$ ) with  $\sum_{i=1}^n p_i = 1$ . If  $f$  is  $m$ -lower convex on  $C$ , then*

$$\frac{m}{2} \sum_{1 \leq i < j \leq n} p_i p_j \|x_i - x_j\|^2 \leq \sum_{i=1}^n p_i f(x_i) - f\left(\sum_{i=1}^n p_i x_i\right).$$

The case of the mappings defined on real intervals have been obtained by Andrica and Raşa in [3].

Furthermore, let us assume that  $\delta(x) := \min_{1 \leq i < j \leq n} \|x_i - x_j\|$ . Then, the following corollary also holds.

**Corollary 2.5** Let  $X, C, f, x_i, p_i$  ( $i = 1, \dots, n$ ) be as in Corollary 2.4. If  $f$  is  $m$ -lower convex on  $C$  with  $m > 0$ , then we have the following refinement of Jensen's inequality:

$$0 < \frac{m}{4} \left( 1 - \sum_{i=1}^n p_i^2 \right) (\delta(x))^2 \leq \sum_{i=1}^n p_i f(x_i) - f \left( \sum_{i=1}^n p_i x_i \right).$$

Counterparts of Proposition 2.3 and Corollaries 2.4, 2.5 for  $M$ -upper convex and  $(m, M)$ -convex functions can be also found in [7].

Now, we will prove the following Jensen's type inequality for strongly convex function with modulus  $c$ .

**Theorem 2.6** Let  $f : C \subseteq X \rightarrow \mathbb{R}$  be a function strongly convex with modulus  $c$  on  $C$  that is open and convex in  $X$ , the normed linear space  $(X, \|\cdot\|)$ ,  $x_i \in C$ ,  $p_i \geq 0$  ( $i = 1, \dots, n$ ) with  $\sum_{i=1}^n p_i = 1$  and  $\bar{x}_p := \sum_{i=1}^n p_i x_i \in C$ . Then, we have the following refinement of Jensen's inequality:

$$\sum_{i=1}^n p_i f(x_i) - f(\bar{x}_p) \geq c \sum_{i=1}^n p_i \|x_i - \bar{x}_p\|^2. \quad (2.3)$$

If  $y \in C$  is such that

$$\sum_{i=1}^n p_i (\nabla_+ f(x_i))(y) \geq \sum_{i=1}^n p_i (\nabla_+ f(x_i))(x_i), \quad (2.4)$$

where  $(\nabla_+ f(\cdot))(\cdot)$  is Gâteaux lateral derivative of  $f$ , then the following refinement of Slater's inequality holds

$$f(y) - c \sum_{i=1}^n p_i \|x_i - y\|^2 \geq \sum_{i=1}^n p_i f(x_i). \quad (2.5)$$

We have the following reverse of Jensen's inequality as well

$$\begin{aligned} & \sum_{i=1}^n p_i (\nabla_+ f(x_i))(x_i) - \sum_{i=1}^n p_i (\nabla_+ f(x_i))(\bar{x}_p) - \sum_{i=1}^n p_i \|x_i - \bar{x}_p\|^2 \\ & \geq \sum_{i=1}^n p_i f(x_i) - f(\bar{x}_p). \end{aligned} \quad (2.6)$$

**Proof** By the definition of  $c$ -strongly convex function on  $C$ , we have

$$t(f(y) - f(x)) \geq f((1-t)x + ty) - f(x) + ct(1-t)\|x - y\|^2$$

for any  $x, y \in C$  and  $t \in [0, 1]$ . This implies that

$$f(y) - f(x) \geq \frac{f(x + t(y - x)) - f(x)}{t} + c(1 - t)\|x - y\|^2$$

for  $t \in (0, 1)$ .

Since  $f$  is convex on open convex subset  $C$ , then the lateral derivative  $(\nabla_+ f(x))(y - x)$  exists for any  $x, y \in C$  and by taking the limit over  $t \rightarrow 0+$ , we get the gradient inequality

$$f(y) - f(x) \geq (\nabla_+ f(x))(y - x) + c\|y - x\|^2 \quad (2.7)$$

for any  $x, y \in C$ .

If we take in (2.7)  $y = x_i, i \in \{1, \dots, n\}$  and  $x = \bar{x}_p$ , then we get

$$f(x_i) - f(\bar{x}_p) \geq (\nabla_+ f(\bar{x}_p))(x_i - \bar{x}_p) + c\|x_i - \bar{x}_p\|^2 \quad (2.8)$$

for any  $i \in \{1, \dots, n\}$ .

Multiply (2.8) by  $p_i \geq 0, i \in \{1, \dots, n\}$  and sum over  $i$  from 1 to  $n$  to get

$$\sum_{i=1}^n p_i f(x_i) - f(\bar{x}_p) \geq \sum_{i=1}^n p_i (\nabla_+ f(\bar{x}_p))(x_i - \bar{x}_p) + c \sum_{i=1}^n p_i \|x_i - \bar{x}_p\|^2. \quad (2.9)$$

This is an inequality of interest in itself.

Since  $(\nabla_+ f(\bar{x}_p))(\cdot)$  is a subadditive and positive homogeneous functional on  $X$ , we have

$$\begin{aligned} \sum_{i=1}^n p_i (\nabla_+ f(\bar{x}_p))(x_i - \bar{x}_p) &\geq (\nabla_+ f(\bar{x}_p))\left(\sum_{i=1}^n p_i x_i - \bar{x}_p\right) \\ &= (\nabla_+ f(\bar{x}_p))(0) = 0 \end{aligned}$$

and by the inequality (2.9), we get the desired result (2.3).

From (2.7), we have for any  $x_i, y \in C, i \in \{1, \dots, n\}$  that

$$f(y) \geq f(x_i) + (\nabla_+ f(x_i))(y - x_i) + c\|x_i - y\|^2$$

for any  $i \in \{1, \dots, n\}$ . If we multiply this inequality by  $p_i \geq 0, i \in \{1, \dots, n\}$  and sum over  $i$  from 1 to  $n$ , then we get

$$f(y) \geq \sum_{i=1}^n p_i f(x_i) + \sum_{i=1}^n p_i (\nabla_+ f(x_i))(y - x_i) + c \sum_{i=1}^n p_i \|x_i - y\|^2. \quad (2.10)$$

By the subadditivity of  $(\nabla_+ f(x_i))(\cdot)$  we have

$$(\nabla_+ f(x_i))(y - x_i) \geq (\nabla_+ f(x_i))(y) - (\nabla_+ f(x_i))(x_i),$$



which implies that

$$\sum_{i=1}^n p_i (\nabla_+ f(x_i))(y - x_i) \geq \sum_{i=1}^n p_i (\nabla_+ f(x_i))(y) - \sum_{i=1}^n p_i (\nabla_+ f(x_i))(x_i),$$

and by (2.10), we get

$$\begin{aligned} f(y) &\geq \sum_{i=1}^n p_i f(x_i) + \sum_{i=1}^n p_i (\nabla_+ f(x_i))(y) - \sum_{i=1}^n p_i (\nabla_+ f(x_i))(x_i) \\ &\quad + c \sum_{i=1}^n p_i \|x_i - y\|^2, \end{aligned} \quad (2.11)$$

for any  $x_i, y \in C$ ,  $i \in \{1, \dots, n\}$  and  $p_i \geq 0$ ,  $i \in \{1, \dots, n\}$  with  $\sum_{i=1}^n p_i = 1$ . This is an inequality of interest in itself.

If the condition (2.4) is valid for some  $y \in C$ , then by (2.11), we get the desired result (2.5).

Now, if we take in (2.11)  $y = \bar{x}_p \in C$ , then we get

$$\begin{aligned} f(\bar{x}_p) &\geq \sum_{i=1}^n p_i f(x_i) + \sum_{i=1}^n p_i (\nabla_+ f(x_i))(\bar{x}_p) - \sum_{i=1}^n p_i (\nabla_+ f(x_i))(x_i) \\ &\quad + c \sum_{i=1}^n p_i \|x_i - \bar{x}_p\|^2 \end{aligned}$$

that is equivalent to the desired result (2.6).  $\square$

**Remark 2.7** For inequalities in terms of the Gâteaux derivatives for convex functions on linear spaces, see [11] while for Slater's type inequalities for convex functions defined on linear spaces and applications, see [12]. The inequalities (2.3)–(2.6) are improvements of the corresponding inequalities for convex functions on normed spaces in which the term  $c \sum_{i=1}^n p_i \|x_i - y\|^2$  vanishes. We observe that, if  $X$  is an inner product space, then using the inner product properties, we have that

$$\sum_{i=1}^n p_i \|x_i - \bar{x}_p\|^2 = \sum_{i=1}^n p_i \|x_i\|^2 - \left\| \sum_{i=1}^n p_i x_i \right\|^2 = \sum_{1 \leq i < j \leq n} p_i p_j \|x_i - x_j\|^2$$

and the inequality (2.2) for  $m > 0$  and the inequality (2.3) for  $c = \frac{m}{2}$  are the same. However, in the general case of normed spaces, they are different. Inequality (2.3) in the case where  $X$  is an inner product space was obtained in [16].

### 3 Some Hermite–Hadamard's Type Inequalities

The classical Hermite–Hadamard double inequality plays an important role in convex analysis and it has a huge literature dealing with its applications, various generalizations and refinements (see for instance [4,9,10,14,20] and the reference therein). In this section, we present Hermite–Hadamard's type inequalities for  $m$ -lower and strongly convex functions.

**Theorem 3.1** *Let  $f : C \subseteq X \rightarrow \mathbb{R}$ , where  $C$  is a convex subset in the normed linear space  $(X, \|\cdot\|)$  and  $x, y \in C$  with  $x \neq y$ . Assume also that  $m > 0$ . If  $f$  is  $m$ -lower convex on  $C$ , then*

$$\begin{aligned} 0 &\leq \frac{1}{16}m \left[ \langle y - x, y + x \rangle_s - \langle y - x, y + x \rangle_i \right] \\ &\leq \frac{1}{2}m \left[ \int_0^1 \|(1-t)x + ty\|^2 dt - \left\| \frac{x+y}{2} \right\|^2 \right] \\ &\leq \int_0^1 f[(1-t)x + ty] dt - f\left(\frac{x+y}{2}\right) \end{aligned} \quad (3.1)$$

and

$$\begin{aligned} 0 &\leq \frac{1}{16}m \left[ \langle y - x, y + x \rangle_s - \langle y - x, y + x \rangle_i \right] \\ &\leq \frac{1}{2}m \left[ \frac{\|x\|^2 + \|y\|^2}{2} - \int_0^1 \|(1-t)x + ty\|^2 dt \right] \\ &\leq \frac{f(x) + f(y)}{2} - \int_0^1 f[(1-t)x + ty] dt. \end{aligned} \quad (3.2)$$

**Proof** The first two inequalities in (3.1) and (3.2) follow by (1.2) and (1.3).

Since  $f$  is  $m$ -lower convex on  $C$ , hence  $g(x) := f(x) - \frac{1}{2}m\|x\|^2$  is convex on  $C$ . Using the Hermite–Hadamard inequality (HH), we have

$$\begin{aligned} f\left(\frac{x+y}{2}\right) - \frac{1}{2}m \left\| \frac{x+y}{2} \right\|^2 &\leq \int_0^1 \left( f[(1-t)x + ty] - \frac{1}{2}m \|(1-t)x + ty\|^2 \right) dt \\ &\leq \frac{1}{2} \left[ f(x) - \frac{1}{2}m\|x\|^2 + f(y) - \frac{1}{2}m\|y\|^2 \right], \end{aligned}$$

which imply the third inequalities in (3.1) and (3.2).  $\square$

**Remark 3.2** If the positivity condition for  $m$  is dropped, then only the third inequalities in (3.1) and (3.2) remain true.

**Corollary 3.3** *Let  $f : C \subseteq X \rightarrow \mathbb{R}$ , where  $C$  is a convex subset in the inner product space  $(X, \langle \cdot, \cdot \rangle)$  and  $x, y \in C$  with  $x \neq y$ . Assume also that  $m > 0$ . If  $f$  is  $m$ -lower*

convex on  $C$ , then

$$\frac{1}{24}m \|x - y\|^2 \leq \int_0^1 f[(1-t)x + ty] dt - f\left(\frac{x+y}{2}\right) \quad (3.3)$$

and

$$\frac{1}{12}m \|x - y\|^2 \leq \frac{f(x) + f(y)}{2} - \int_0^1 f[(1-t)x + ty] dt. \quad (3.4)$$

**Proof** Since  $(X, \langle \cdot, \cdot \rangle)$  is an inner product, then for any  $x, y$

$$\begin{aligned} & \int_0^1 \|(1-t)x + ty\|^2 dt \\ &= \int_0^1 \left[ (1-t)^2 \|x\|^2 + 2t(1-t) \operatorname{Re} \langle x, y \rangle + t^2 \|y\|^2 \right] dt \\ &= \|x\|^2 \int_0^1 (1-t)^2 dt + 2\operatorname{Re} \langle x, y \rangle \int_0^1 t(1-t) dt + \|y\|^2 \int_0^1 t^2 dt \\ &= \frac{1}{3} \left( \|x\|^2 + \operatorname{Re} \langle x, y \rangle + \|y\|^2 \right). \end{aligned}$$

Therefore, for any  $x, y$

$$\begin{aligned} & \int_0^1 \|(1-t)x + ty\|^2 dt - \left\| \frac{x+y}{2} \right\|^2 \\ &= \frac{1}{3} \left( \|x\|^2 + \operatorname{Re} \langle x, y \rangle + \|y\|^2 \right) - \frac{1}{4} \left( \|x\|^2 + 2\operatorname{Re} \langle x, y \rangle + \|y\|^2 \right) \\ &= \frac{1}{12} \left( \|x\|^2 - 2\operatorname{Re} \langle x, y \rangle + \|y\|^2 \right) = \frac{1}{12} \|x - y\|^2 \end{aligned}$$

and

$$\begin{aligned} & \frac{\|x\|^2 + \|y\|^2}{2} - \int_0^1 \|(1-t)x + ty\|^2 dt \\ &= \frac{1}{2} \left( \|x\|^2 + \|y\|^2 \right) - \frac{1}{3} \left( \|x\|^2 + \operatorname{Re} \langle x, y \rangle + \|y\|^2 \right) \\ &= \frac{1}{6} \|x - y\|^2. \end{aligned}$$

Using Theorem 3.1, we get the desired results (3.3)–(3.4).  $\square$

**Remark 3.4** If  $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$  is twice differentiable and  $f''(t) \geq m$  for any  $t \in [a, b]$ , then by (3.3) and (3.4), we have

$$\frac{1}{24}m(b-a)^2 \leq \frac{1}{b-a} \int_a^b f(s) ds - f\left(\frac{a+b}{2}\right), \text{ see [8, Eq. (5.1)]}$$

and

$$\frac{1}{12}m(b-a)^2 \leq \frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(s) ds, \text{ see [13],[14, p. 40].}$$

The next theorem gives a counterpart of the Hermite–Hadamard inequality for strongly convex functions.

**Theorem 3.5** *Let  $f : C \subseteq X \rightarrow \mathbb{R}$  be a function strongly convex with modulus  $c$  on  $C$  that is open and convex in the normed linear space  $(X, \|\cdot\|)$  and  $x, y \in C$  with  $x \neq y$ . Then*

$$\frac{f(x) + f(y)}{2} - \int_0^1 f[(1-t)x + ty] dt \geq \frac{1}{6}c \|x - y\|^2 \quad (3.5)$$

and

$$\int_0^1 f[(1-t)x + ty] dt - f\left(\frac{x+y}{2}\right) \geq \frac{1}{12}c \|x - y\|^2. \quad (3.6)$$

**Proof** If we integrate condition (2.1) in the definition of strongly convex functions, we have

$$\begin{aligned} \frac{f(x) + f(y)}{2} - \int_0^1 f(tx + (1-t)y) dt &\geq c \|x - y\|^2 \int_0^1 t(1-t) dt \\ &= \frac{1}{6}c \|x - y\|^2 \end{aligned}$$

for any  $x, y \in C$ , which proves (3.5).

By taking in the definition of strong convexity  $t = \frac{1}{2}$ , we have

$$\frac{f(a) + f(b)}{2} - f\left(\frac{a+b}{2}\right) \geq \frac{c}{4} \|a - b\|^2, \quad (3.7)$$

for all  $a, b \in C$ .

If we take in (3.7)  $a = (1-t)x + ty$  and  $b = tx + (1-t)y$ , then we get

$$\frac{f((1-t)x + ty) + f(tx + (1-t)y)}{2} - f\left(\frac{x+y}{2}\right) \geq c \left(t - \frac{1}{2}\right)^2 \|x - y\|^2$$

for any  $x, y \in C$  and  $t \in [0, 1]$ .

Integrating over  $t \in [0, 1]$ , we have

$$\begin{aligned} &\frac{1}{2} \left[ \int_0^1 f((1-t)x + ty) dt + \int_0^1 f(tx + (1-t)y) dt \right] - f\left(\frac{x+y}{2}\right) \\ &\geq c \|x - y\|^2 \int_0^1 \left(t - \frac{1}{2}\right)^2 dt \end{aligned} \quad (3.8)$$

and since

$$\int_0^1 f((1-t)x + ty) dt = \int_0^1 f(tx + (1-t)y) dt \text{ and } \int_0^1 \left(t - \frac{1}{2}\right)^2 dt = \frac{1}{12},$$

then by (3.8), we get (3.6).  $\square$

**Remark 3.6** If  $f : [a, b] \subset \mathbb{R} \rightarrow \mathbb{R}$  is a function strongly convex with modulus  $c$  on the interval  $[a, b]$ , then by (3.5) and (3.6), we get

$$\frac{f(a) + f(b)}{2} - \frac{1}{b-a} \int_a^b f(s) ds \geq \frac{1}{6} c (b-a)^2$$

and

$$\frac{1}{b-a} \int_a^b f(s) ds - f\left(\frac{a+b}{2}\right) \geq \frac{1}{12} c (b-a)^2.$$

These inequalities were obtained in this form in [16].

**Open Access** This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

## References

- Adamek, M.: On a problem connected with strongly convex functions. *Math. Inequal. Appl.* **19**, 1287–1293 (2016)
- Abbaszadeh, S., Gordji, M., Eshaghi: On the quadratic support of strongly convex functions. *Int. J. Nonlinear Anal. Appl.* **7**(1), 15–20 (2016)
- Andrica, D., Raşa, I.: The Jensen inequality: refinements and applications. *Anal. Numer. Theor. Approx.* **74**, 105–108 (1985)
- Bessenyei, M., Páles, Z.S.: Characterization of convexity via Hadamard's inequality. *Math. Inequal. Appl.* **9**, 53–62 (2006)
- Bakula, M., Klaričić, Nikodem, K.: On the converse Jensen inequality for strongly convex functions. *J. Math. Anal. Appl.* **434**, 516–522 (2016)
- Cioranescu, I.: *Geometry of Banach Spaces, Duality Mappings and Nonlinear Problems*. Kluwer Academic Publishers, Dordrecht (1990)
- Dragomir, S.S.: Some inequalities for  $(m, M)$ -convex mappings and applications for the Csiszár  $\Phi$ -divergence in information theory. *Math. J. Ibaraki Univ.* **33**, 35–50 (2001)
- Dragomir, S.S.: On the Jensen's inequality for isotonic linear functionals. *Nonlinear Anal. Forum* **7**(2), 139–151 (2002)
- Dragomir, S.S.: An inequality improving the first Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products. *J. Inequal. Pure Appl. Math.* **3**(2), 8 (2002). (Article 31)
- Dragomir, S.S.: An inequality improving the second Hermite-Hadamard inequality for convex functions defined on linear spaces and applications for semi-inner products. *J. Inequal. Pure Appl. Math.* **3**(3), 8 (2002). (Article 35)
- Dragomir, S.S.: Inequalities in terms of the Gâteaux derivatives for convex functions on linear spaces with applications. *Bull. Aust. Math. Soc.* **83**(3), 500–517 (2011)

12. Dragomir, S.S.: Some Slater's type inequalities for convex functions defined on linear spaces and applications. *Abstr. Appl. Anal.* **2012**, 16 (2012). (**Art. ID 168405**)
13. Dragomir, S.S., Cerone, P., Sofo, A.: Some remarks on the trapezoid rule in numerical integration. *Indian J. Pure Appl. Math.* **31**(5), 475–494 (2000)
14. Dragomir, S.S., Pearce, C.E.M.: *Selected Topics on Hermite–Hadamard Inequalities and Applications*. RGMIA Monographs, Victoria University (2000). <http://rgmia.org/monographs.php>. Accessed 20 June 2018
15. Hiriart-Urruty, J.B., Lemaréchal, C.: *Fundamentals of Convex Analysis*. Springer-Verlag, Berlin Heidelberg (2001)
16. Merentes, N., Nikodem, K.: Remarks on strongly convex functions. *Aequ. Math.* **80**, 193–199 (2010)
17. Merentes, N., Nikodem, K.: Strong convexity and separation theorems. *Aequ. Math.* **90**, 47–55 (2016)
18. Montruccio, L.: Lipschitz continuous policy functions for strongly concave optimization problems. *J. Math. Econ.* **16**, 259–273 (1987)
19. Moradi, H.R., Omidvar, M.E., Khan, M.A., Nikodem, K.: Around Jensen's inequality for strongly convex functions. *Aequat. Math.* **92** (2018), 25–37. <https://doi.org/10.1007/s00010-017-0496-5>
20. Niculescu, C.P., Persson, L.-E.: *Convex Functions and their Applications. A Contemporary Approach*. CMS Books in Mathematics, vol. 23. Springer-Verlag, New York (2006)
21. Nikodem, K.: On strongly convex functions and related classes of functions. In: Rassias, T.M. (ed.) *Handbook of Functional Equations. Functional Inequalities*, 365–405, Springer Optimization and Its Application 95. Springer, New York (2015)
22. Nikodem, K., Páles, Z.S.: Characterizations of inner product spaces by strongly convex functions. *Banach J. Math. Anal.* **5**(1), 83–87 (2011)
23. Polyak, B.T.: Existence theorems and convergence of minimizing sequences in extremum problems with restrictions. *Soviet Math. Dokl.* **7**, 72–75 (1966)
24. Rajba, T., Wąsowicz, S.Z.: Probabilistic characterization of strong convexity. *Opuscula Math.* **31**(1), 97–103 (2011)
25. Roberts, A.W., Varberg, D.E.: *Convex Functions*. Academic Press, New York, London (1973)
26. Vial, J.P.: Strong convexity of sets and functions. *J. Math. Econ.* **9**, 187–205 (1982)