



VICTORIA UNIVERSITY
MELBOURNE AUSTRALIA

Development and psychometric validation of the mental health-related barriers and benefits to EXercise (MEX) scale in healthy adults

This is the Published version of the following publication

Connolly, Madeleine, Bowden, Stephen C, Pascoe, Michaela and Van Dam, Nicholas T (2023) Development and psychometric validation of the mental health-related barriers and benefits to EXercise (MEX) scale in healthy adults. *Sports Medicine - Open*, 9 (18). ISSN 2199-1170

The publisher's official version can be found at
<https://sportsmedicine-open.springeropen.com/articles/10.1186/s40798-023-00555-x>
Note that access to this version may require subscription.

Downloaded from VU Research Repository <https://vuir.vu.edu.au/46960/>

ORIGINAL RESEARCH ARTICLE

Open Access



Development and Psychometric Validation of the Mental Health-Related Barriers and Benefits to EXercise (MEX) Scale in Healthy Adults

Madeleine L. Connolly^{1*} , Stephen C. Bowden^{1,2}, Michaela C. Pascoe³ and Nicholas T. Van Dam¹

Abstract

Background Physical exercise has been shown to reduce anxiety and depression symptoms, the most common mental health disorders globally. Despite the benefits of exercise in anxiety and depression, the symptoms of these disorders may directly contribute to a lack of engagement with exercise. However, mental health-related barriers and benefits to exercise engagement have not been addressed in quantitative research. We introduce the development and psychometric validation of the Mental health-related barriers and benefits to EXercise (MEX) scale.

Methods Three samples were collected online prospectively (sample 1 $n=492$; sample 2 $n=302$; sample 3 $n=303$) for scale refinement and validation with exploratory and confirmatory factor analysis. All participants were generally healthy adults, aged 18–45, and had no history of severe mental illness requiring hospitalization and no physical disability impacting over 50% of daily function.

Results We identified a 30-item, two-factor model comprising 15 barrier and 15 benefit items. Overall model fit was excellent for an item-level scale across the three samples (Comparative Fit Index = 0.935–0.951; Root-Mean-Square Error of Approximation = 0.037–0.039). Internal consistency was also excellent across the three samples ($\alpha=0.900$ –0.951). The barriers subscale was positively correlated with symptoms of anxiety, depression and stress, and negatively correlated with measures of physical activity and exercise engagement. The benefits subscale was negatively correlated with symptoms of anxiety, depression and stress, and positively correlated with measures of physical activity and exercise engagement.

Conclusion The MEX is a novel, psychometrically robust scale, which is appropriate for research and for clinical use to ascertain individual and/or group level mental health-related barriers and benefits to exercise.

Keywords Anxiety, Depression, Stress, Barriers, Benefits, Exercise

Background

Anxious and depressive disorders are the most common mental disorders among adults worldwide and leading causes of disability [1]. As a result, the promotion of mental health and the prevention of mental health disorders is paramount. Exercise, defined here as effortful physical activity in leisure time, is a promising approach for the prevention and treatment for symptoms of both anxiety and depression at both clinical and subclinical levels [2].

*Correspondence:

Madeleine L. Connolly

mconnolly@student.unimelb.edu.au

¹ Melbourne School of Psychological Sciences, The University of Melbourne, Level 12, Redmond Barry Building, Parkville, Melbourne 3010, Australia

² Centre for Clinical Neurosciences and Neurological Research, St Vincent's Hospital Melbourne, Melbourne 3065, Australia

³ Institute for Health and Sport, Victoria University, Melbourne 3011, Australia



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

As a treatment option, exercise requires motivation and action for uptake and adherence. However, factors such as a pervasive lack of motivation and fear-based avoidance—key clinical features of depression and anxiety, respectively—may potentially create barriers to engagement in leisure-time exercise for individuals who experience such symptoms [3, 4]. Other symptoms, such as concerns about appearance, sensitivity to somatic signals (e.g. heart rate, respiration), and avoidance, may also critically interact with uptake of and adherence to exercise. Furthermore, there are broader psychological benefits to exercise beyond the overall score reduction on anxiety or depressive scales that are typically reported for clinical [5] and subclinical populations [2]. Thus, an important bidirectional relationship may apply. While exercise may reduce or improve specific symptoms of anxiety and depression, the experience of the symptoms themselves may prevent people from exercising.

Measurement of mental health-related barriers to, and benefits of exercise engagement is crucial to understanding the bidirectional relationship and, for achieving appropriate, tailored exercise prescriptions in individuals with symptoms of anxiety and depression. It is important to acknowledge a distinction between exercise and physical activity (PA). Whilst exercise refers to effortful physical activity in leisure time specifically, PA refers to all activity across life domains including leisure time, but also including any movement associated with occupation, transport, housework, and any other non-leisure-time domains. Although many types of PA are beneficial for physical health, leisure-time exercise is particularly beneficial for mental health [6]. It is also important to acknowledge both the barriers and benefits to exercise engagement during early adulthood, as this period is marked by a decline in PA and exercise engagement [7]. Furthermore, early adulthood is also a common time for the onset of mental disorders, with recent estimates of median ages of onset of anxiety and mood disorders being 17 and 31, respectively [8].

Current quantitative models of benefits and barriers to exercise are based on scales that are intended for the general adult population and cover a wide range of general, logistical and physical barriers and benefits to exercise, without a distinct focus on individual mental health [9]. Consequently, these models are missing specific features of anxiety- and depression-like symptoms that may be relevant to exercise behaviours, in both clinical and subclinical populations. In essence, quantitative research into the barriers and benefits to exercise engagement only assesses a small subset of psychologically relevant items, always combined with a much larger list of physical or logistical barriers and benefits to exercise [3, 9–11]. Meanwhile, qualitative research has demonstrated that

the actual experience of anxious and depressive symptoms appear to become barriers to exercise engagement, whilst the relief of these symptoms is a benefit of exercise engagement [12, 13]. Despite such qualitative evidence, specific quantitative assessment of mental health-related barriers and benefits to exercise is critical to understanding how symptoms are related to exercise.

Mental Health-Related Benefits of Exercise

In their meta-analysis of the benefits and barriers to exercise for depressive and mood disorders, Glowacki et al. report improved mood, better attention and alertness, and better stress management as major benefits from exercise [3]. Similarly, Searle and colleagues' qualitative evidence from a sample of 33 depressed individuals suggests that positive distraction from depressive symptoms and providing a sense of purpose are key benefits of exercise [4]. Improvements in energy, a positive feeling of 'pushing oneself', a sense of regaining pleasure, and increases in self-confidence, self-efficacy, body image, and autonomy are also factors reported by participants with depression and following exercise engagement in qualitative research [4, 13]. Accordingly, several key symptoms of depression (including low mood, high stress, low energy, anhedonia, low self-confidence and low self-efficacy) are improved with exercise.

Qualitative evidence also points to psychologically relevant benefits of exercise for those with anxiety disorders, relating to both the alleviation of anxious symptoms and of general negative affect. Mason et al. [12] interviewed a heterogeneous sample of individuals with symptoms of at least one anxiety or related disorder (e.g. obsessive-compulsive disorder, specific phobia). Following exercise engagement, they observed decreases in frequency and severity of worry (especially among those with generalized anxiety), social benefits (particularly by those with social anxiety symptoms), and the perception that exercise was a useful distraction from anxious thoughts. Finally, exercise was also reported to reduce overall stress levels among individuals with anxiety and depression [3, 4]. It is important to note that there are many psychological and mental health-related benefits of exercise, and the benefits reported here are not exhaustive. Further information on mental health-related benefits of exercise is available in Mikkelsen et al.'s review [14].

Mental Health-Related Barriers to Exercise

Barriers to exercise which mirror depressive and anxious symptoms appear frequently in qualitative data from clinical samples, and to a smaller extent in quantitative data representing subclinical samples. Such barriers are not only limited to the more commonly cited barriers of lack of motivation and fear-based avoidance, but also

include a variety of transdiagnostic mental health-related barriers to exercise [3, 4].

In quantitative research for non-mental health specific barriers to exercise, some of the most highly endorsed barriers in the general population samples were: low confidence when sad or distressed (75% endorsement in a sample of 138), fatigue (69%, total sample $n=203$), low mood (65%, total sample $n=102$), feeling unwell (60%, total sample $n=101$), and feeling too shy or embarrassed (36%, total sample $n=101$) [3]. Interestingly, of these five barriers, three (low confidence, fatigue, and low mood) directly mirror symptoms of depression, and one (feeling too shy or embarrassed) mirrors social anxiety. Similarly, in qualitative research among individuals diagnosed with depressive disorders, there are many reports of barriers to exercise which mirror depressive symptoms, such as low motivation and low confidence [3, 4]. The qualitative research also points to further barriers to exercise in clinical depression samples that are more like anxious or transdiagnostic symptoms, such as guilt, stress, fear of social interaction, and poor body image [4].

Exercise-related anxiety (i.e. anxiety specific to the activity of exercising) has also been reported as a barrier to exercise. For individuals with high levels of symptomatic anxiety, qualitative interviews indicate that specific fears or anxieties can negatively affect exercise participation and become barriers to exercise. For example, participants with social anxiety report fearing social evaluation during exercise, and consequently avoid exercise behaviours [12]. For participants with general anxiety, fears such as injury or pain were also key barriers. Additionally, psychosomatic sensations of anxiety during exercise were also reported as barriers to future exercise sessions by some with anxiety [12]. Transdiagnostic symptoms of low self-confidence, fatigue, life stress, and worries of lack of exercise knowledge or technique have also been noted as barriers to exercise for individuals with high levels of anxious symptoms [12].

As with the benefits of exercise, the mental health-related barriers to exercise reported by individuals with depressive and anxious symptoms show considerable transdiagnostic overlap. Such barriers can be labelled as low confidence, low mood, fatigue, stress, body image, worry, social anxiety, fear of injury, and lack of exercise knowledge and skills.

The Novel Mental Health-Related Barriers and Benefits to Exercise Scale

The aim of this study was to develop and validate a novel scale with items specific to the barriers and benefits of physical exercise, relevant to mental health and psychological well-being. The scale was intended to assess the mental health-related barriers and benefits of exercise for

individuals with both varied levels of anxious and depressive symptoms. The scale therefore aimed to quantify the psychological barriers and benefits to exercise in generally healthy adults without severe mental illness.

It was hypothesized that the *Mental health-related barriers and benefits to physical EXercise* (MEX) scale would comprise two main factors, a barriers subscale and a benefits subscale. Within each subscale, multiple lower-order factors were explored.

It was also hypothesized that the two subscales would have associations with measures of mental health, psychological well-being, and physical activity and exercise. The barriers subscale was predicted to have positive correlations with mental ill health, and negative correlations with psychological well-being, and with physical activity and exercise. Meanwhile, the benefits subscale was predicted to have a negative correlation with mental ill health, but positive correlations with psychological well-being, and with physical activity and exercise.

Methods

Participants and Design

The present study employed data from two data sets, the second of which was randomly divided, to create three samples in total. Recruitment of all participants was aimed at the general population, with the aim of recruiting generally healthy adults without severe mental illness. Sample 1, the initial scale development data set, was procured using Amazon's Mechanical Turk (Mturk) in 2020 (final $n=492$). The second data set consisted of participants recruited via Prolific in 2021 (final $n=605$). This second data set was split (sample 2: $n=302$; sample 3: $n=303$) so that Sample 2 data could be used in scale refinement, and Sample 3 data remained a separate validation sample. Given the difficulty of ascertaining power for factor analysis [15], we aimed to recruit as many participants as possible in both stages of recruitment.

In the first data set, geographical location was restricted to US-based participants to mitigate potential confounds associated with geo-cultural differences in item interpretation. In the second data set, geographical location was restricted to residents of the USA, the UK, Australia, and New Zealand; representation was expanded to allow for greater ecological validity. All participants across each data set were between 18 and 45 years of age (to minimize age-related variation in activity level), and were deemed eligible on the basis of satisfying the following inclusion criteria: (1) no reported history of severe mental illness; (2) no self-reported physical disability perceived by participant to negatively impact >50% of their daily function. Participants from each data set were also excluded during data cleaning processes if they did not satisfy the attention check and careful responding criteria

as listed below (see *Data Cleaning Procedure*). Full demographic information and data processing outcomes are described under *Results*. Participants were asked about their experience of mental health symptoms, as well as their current activity levels. Upon completion of the total battery, participants were remunerated at a minimum rate of \$10US per hour. The collection of data for both data sets was approved by the University of Melbourne Human Research Ethics Committee [HREC # 2056662], and all protocols conformed to the ethical standards outlined in the Declaration of Helsinki.

Measures

The MEX Scale

The initial pool numbered 113 items (53 benefits, 58 barriers, 1 item for 'benefit facilitation', 1 item for 'barrier interference'). A number of items for the Barrier and Benefits subscales in the MEX were inspired by psychologically relevant barrier and benefit items from prior quantitative research [9–11, 16–19]. Further, novel quantitative items were written by the authors with inspiration from qualitative research on depression [13], and on anxiety disorders [12]. Finally, a small number of novel items were also written by the authors with inspiration from informal piloting with 30 adults from the general population, and from the authors' clinical knowledge and expertise. All 113 items were administered to sample 1. A subset of the MEX scale (48 items; 16 benefit items and 32 barrier items) was administered to sample 2/3.

Depression, Anxiety, and Stress Scales (DASS-42)

The DASS-42 is a 42-item scale designed to measure depression, anxiety, and stress on a continuum [20]. The DASS-42 has been validated previously by factor analysis and shows good reliability [20]. Each of the DASS-42 items is scored on a 4-point scale, in which higher scores indicate higher levels of depression, anxiety, or stress [20]. The DASS-42 was only administered to Sample 1. Each subscale showed a bimodal distribution in Sample 1.

The International Physical Activity Questionnaire—Short Form (IPAQ-SF)

The IPAQ-SF is a 7-item questionnaire designed to measure the weekly minutes of all physical activity and transform these activity minutes into metabolic equivalent units (METs) [21]. The IPAQ-SF has participants recall and report weekly hours and/or minutes of all physical activity, including work and home domains, as well as leisure-time activity. Participants reported the activity as either 'vigorous', 'moderate', or 'walking'. Each category of intensity was then multiplied by the corresponding metabolic equivalent and then summed for a score of METs

over one week, where higher METs indicate higher levels of activity. The IPAQ-SF was chosen here for its brevity in administration; however, it has previously shown weak external validity when correlated with objective accelerometer measures, and self-report appears to be more prone to overestimation by those who are more sedentary [22]. The IPAQ-SF was only administered to Sample 1.

The Satisfaction With Life Scale (The SWLS)

The SWLS is a 5-item scale, designed to measure overall satisfaction with life [23]. The SWLS has been validated previously by factor analysis, and shows good external validity [23]. Each of the SWLS items is scored on a 6-point scale, in which higher scores indicate higher levels of life satisfaction and psychological well-being [23]. The measure was chosen as a counterbalance for the DASS-42, which only measures negative affect. The SWLS was only administered to Sample 1.

The Balanced Inventory of Desirable Responding Scale—Short Form (The BIDR)

The BIDR is a 16-item scale designed to measure the extent to which participants have bias in responses either in terms of a) impression management (8 items), or b) self-deceptive enhancement (8 items) [24]. The BIDR has been validated previously by factor analysis, and shows good reliability [24]. Each of the BIDR items is scored on a 5-point scale, in which higher scores indicate higher levels of either impression management or self-deceptive enhancement [24]. This scale was selected to assess the extent to which the MEX subscales would correlate with either measure of response bias. The BIDR was only administered to Sample 1.

Subjective Activity and Fitness Questions

All participants were presented with questions regarding how they would evaluate their own physical activity and fitness in terms of general sufficiency, in terms of their ideal levels of each, and each as compared to the average person of their age [25]. Such questions have previously shown high external validity and correlations with psychological measures such as distress and sleep [25]. Higher scores on subjective activity and fitness indicate higher perceived levels of activity and fitness. These questions were administered to both samples.

COVID-19 Impact Questions

All participants were asked questions to assess the potential negative impact and extent of any impact of the COVID-19 pandemic on their mental health, and exercise habits. These questions were administered to both samples.

The Mood and Anxiety Symptom Questionnaire-90 (MASQ-90)

The MASQ-90 is a 90-item questionnaire designed to measure: Anxious Arousal, Anhedonic Depression, and three types of Symptoms of General Distress: Depressive Symptoms, Anxious Symptoms, and Mixed Symptoms [26]. The MASQ-90 has been validated previously by factor analysis, and shows good reliability [26]. Each of the MASQ-90 items is scored on a 5-point scale, in which higher scores indicate higher levels of depression, anxiety, or distress [26]. The MASQ-90 was only administered to Sample 2/3. Each of the subscales was positively skewed in distribution for samples 2/3.

Leisure Time Sport and Exercise Survey (LTSES)

The LTSES is a brief scale designed to capture habitual physical activity across the domain of sport and exercise activity in leisure time [27]. Participants are asked to report their two most frequented sports or leisure-time activities (e.g. running, walking, or gym-based activity) and to report the intensity of each activity, and the time spent on each activity in terms of months per year and minutes per week. The overall score is transformed to a number between 0 and 4, in which 0 corresponds to no leisure-time exercise and 4 corresponds to a very high amount of leisure-time exercise [27]. The LTSES has been validated previously by factor analysis and shows good reliability [27]. The LTSES was only administered to Sample 2/3.

Data Cleaning Procedure

Data were excluded casewise if it did not satisfy the following criteria: a recaptcha score of over 0.7 (Qualtrics recommended threshold), and passing all attention checks and basic writing and reading comprehension checks. The data were also screened for long string response outliers [28], high intra-individual response variability (IRV) [29], and multivariate outliers by Mahalanobis distance. Long string response outliers were judged by visual inspection of the long string response histogram per data set. High IRV was determined by the score corresponding to the 99th percentile of IRV scores in a randomized simulation of each data set. Finally, the Mahalanobis distance outlier cut-off score was determined by the value corresponding to the 99.9th percentile of the chi-square distribution of Mahalanobis distances for the remaining participants in each sample after prior cases had been removed. Full details of data cleaning results for each sample are available in Additional file 1.

Data Analysis Procedure

Data were analysed in R (version 4.1.0). For exploratory factor analysis (EFA), we used promax rotation and maximum likelihood estimation. For confirmatory factor analysis (CFA), we used weighted least squares mean and variance adjusted estimation, due to its robustness with ordered categorical items. Fit statistics were judged against the Hu and Bentler criteria for good fit: Tucker–Lewis Index (TLI) and Comparative Fit Index (CFI) above 0.95, Standardized Root-Mean-Squared Residual (SRMR) under 0.08, and Root-Mean-Square Error of Approximation (RMSEA) under 0.06 [30]. We note that these are general and conservative heuristics, which are better for parcel-level factor analysis; we therefore apply some leniency in the interpretation of our item-level analysis [31, 32].

Following factor analytic procedures, the reliability of the scale was determined using both alpha and omega coefficients. Reliability for each alpha and omega was judged against the traditional criteria of a threshold of 0.7 or above meaning good fit for each coefficient, and 0.9 or above meaning excellent fit [32]. The convergent validity of the barriers and benefits scales was examined via correlational estimates of each subscale to external variables. For each pair of correlations, a Fisher's Z score was also calculated to examine relative strength of correlations between the external measure and the barriers scale versus the external measure with the benefits scale. The Fisher's Z scores were calculated with the Hittner et al. modification of Dunn and Clark's procedure to adjust for correlated correlations [33].

Results

Demographics

The first sample ($n=492$) was largely male (65.85%), Caucasian (70.2%), college educated (73%), employed full time (80.4%), and in their 30s ($M=32.4$, $SD=5.69$). The second and third samples comprise random split halves of a larger sample ($n=605$). The second sample ($n=302$) was largely female (79.1%), Caucasian (55%), high school educated (56.7%), and in their 20s ($M=24.1$, $SD=6.5$). Similarly, the third sample ($n=303$) was largely female (80.2%), Caucasian (56.8%), high school educated (58%), and in their 20s ($M=23.8$, $SD=6.1$). A summary of demographic statistics are given in Table 1; full demographic statistics are shown in Additional file 1: Table S1.

Item Reduction and Final Scale

Parallel Analysis and initial EFA: Sample 1

Parallel analysis [34] of the initial 112 items suggested four factors. The ensuing four factor EFA model was poorly specified, with no primary loadings on the fourth

Table 1 Brief demographic statistics for all Samples 1, 2, and 3

	Sample 1 (n = 492)	Sample 2 (n = 302)	Sample 3 (n = 303)
	M (SD)	M (SD)	M (SD)
Age	32.4 (5.69) % (n = 492)	24.1 (6.5) % (n = 302)	23.8 (6.1) % (n = 303)
Gender (%)			
Male	65.85%	19.87%	18.48%
Female	34.15%	79.14%	80.2%
Non-binary	0%	0.99%	1.32%
Country of residence (%)			
The USA	100%	83.11%	82.84%
The UK	0%	3.97%	2.97%
Australia	0%	7.95%	11.22%
New Zealand	0%	4.97%	2.97%
Race (%)			
White	70.33%	54.97%	56.77%
Hispanic or Latino	5.28%	14.24%	14.52%
Asian	5.69%	14.57%	14.52%
Native American, Alaskan Native, Native Hawaiian or Pacific Islander	0.61%	0.99%	0.99%
Black or African American	16.46%	9.6%	6.27%
Multiple Races/Unknown/Do Not Wish to Disclose	1.63%	5.63%	6.93%

Full demographic statistics are provided in Additional file 1: Table S1

factor. A second EFA was estimated with two factors, to explore the theoretical assumption of one benefits factor and one barriers factor. The two-factor model item loadings reflected one clear ‘benefits’ factor and one clear ‘barriers’ factor. Despite a coherent factor structure, the two-factor EFA model showed poor overall fit with TLI = 0.78. Given the poor overall fit, and the difficulty of interpreting 112 items, item reduction was undertaken.

Refinement EFAs and CFAs: Samples 1 and 2

Items were removed in an iterative process considering the following issues: i) high cross-loadings in EFA (cross-loading above a magnitude of 0.2 on multiple factors); ii) loadings below 0.3 in the specified factor in CFA; iii) poor theoretical fit of items.

A resulting two-factor EFA model with 46 items showed a robust factor structure, with factor 1 a ‘barriers’ factor, and factor 2 a ‘benefits’ factor, and no high cross-loadings. The chi-square goodness-of-fit statistic was significant ($\chi^2 = 2281.34$, $df = 944$, $p < 0.001$); however, overall fit was adequate in Sample 1, with a TLI of 0.89, an RMSEA index of 0.054 (90% CI: lower = 0.051, upper = 0.057) and SRMR estimate of 0.04.

A two-factor CFA was estimated based on the prior EFA. The two-factor CFA model showed all 46 items loading significantly. The CFA model showed good overall fit on all indices, except for a significant chi-square

goodness of fit: $\chi^2 (988) = 1649.1$, $p < 0.001$. The CFI of 0.95, TLI of 0.94, RMSEA of 0.37, and SRMR estimate of 0.062 all showed good fit.

The two-factor 46-item model of the MEX had poor fit in Sample 2 based on a CFI of 0.83, and a TLI of 0.83, prompting further revision. Iterative EFA and CFA estimations were completed after removing items causing poor model-to-data fit in the second sample, resulting in 15 barrier items, and 15 benefit items as the final MEX scale. Two-factor confirmatory factor analysis of the final MEX scale showed good model fit overall in Sample 2, despite a significant chi-square goodness of fit (see Table 3). All items loaded significantly on their respective factors (see Table 2). The inter-factor correlation was negative and moderate in Sample 2, $r = -0.479$.

The refined 30-item, two-factor MEX showed good model fit in Sample 1, despite a significant chi-square goodness of fit (see Table 3). All primary item loadings were significant (see Table 2). Inter-factor correlations were weak and negative in Sample 1, $r = -0.11$.

Confirmation CFA: Sample 3

A two-factor CFA of the 30-item MEX showed overall good fit in the third sample, despite a significant chi-square goodness of fit (see Table 3). All primary item loadings were significant (see Table 2). Inter-factor correlations were moderate and negative, $r = -0.49$.

Table 2 Standardized CFA loadings across the three samples on the 30-item MEX two-factor model

Factor	Items	Sample 1 (n = 492)	Sample 2 (n = 302)	Sample 3 (n = 303)	
Benefits	Exercise helps me to have a positive outlook on life	0.549	0.714	0.711	
	Exercise makes me feel more self-confident	0.614	0.559	0.672	
	Exercising gives me more energy	0.634	0.683	0.783	
	Exercise helps me to manage my stress	0.673	0.805	0.795	
	Exercise benefits my mental health	0.605	0.722	0.786	
	Exercise helps me to feel less angry or irritable	0.588	0.727	0.645	
	After exercising, I notice a decrease in my general level of worry and concern	0.588	0.681	0.670	
	Regular exercise makes feel me more mentally alert	0.589	0.710	0.704	
	Exercise helps me to feel motivated again	0.602	0.626	0.696	
	Exercise gives me an instant mood increase	0.598	0.720	0.743	
	Exercise makes me feel more energetic	0.651	0.731	0.694	
	Exercise helps me to cope with life's stresses	0.660	0.768	0.813	
	Exercise helps me manage my mood	0.626	0.782	0.761	
	Exercising changes my attitude	0.633	0.668	0.668	
	Exercise gives me a long term mood increase	0.521	0.665	0.708	
	Barriers	I'm too shy or embarrassed to start exercising	0.803	0.759	0.798
		I feel self-conscious about exercising	0.656	0.704	0.762
		I'm worried exercise will hurt	0.749	0.446	0.497
I'm not confident in my ability to exercise		0.784	0.795	0.808	
Mental exhaustion stops me from exercising		0.762	0.787	0.798	
I worry that I would not be very good at exercising		0.775	0.649	0.618	
Exercising can trigger feelings of anxiety for me		0.786	0.733	0.800	
I feel embarrassed if people see me exercising		0.785	0.486	0.498	
Some days I can't get out of bed, let alone exercise		0.802	0.457	0.425	
I don't want to exercise because I don't like how my body looks		0.750	0.656	0.657	
I don't want to exercise when I am angry, or in a bad mood		0.721	0.509	0.438	
I worry about what other people think of me exercising		0.727	0.69	0.764	
My exercise or fitness goals seem too hard to achieve		0.662	0.567	0.571	
I've failed to achieve too many exercise goals in the past		0.775	0.614	0.515	
I'm scared of losing my breath while exercising		0.731	0.491	0.513	

All loadings statistically significant, $p < .05$

Table 3 CFA fit statistics for the two-factor model of the final 30-item MEX scale in Sample 1, 2, and 3

	CFI	TLI	X2	RMSEA (point)	RMSEA 90% CI (lower)	RMSEA 90% CI (upper)	SRMR
Sample 1	0.951	0.947	679.30*	0.037	0.032	0.042	0.052
Sample 2	0.946	0.942	584.27*	0.039	0.031	0.045	0.056
Sample 3	0.935	0.930	590.36*	0.039	0.032	0.046	0.054

Statistics are robust estimates.

* $p < .001$, $df = 988$

The final CFA overall fit statistics for each sample can be seen in Table 3. Each row in Table 3 corresponds to the fit statistics of the 2-Factor CFA models in the three samples. Each 2-factor model comprises the 30 final MEX items, as two 15-item factors (benefits and barriers). The fit statistics listed are robust estimates.

Descriptive Statistics, Reliability, and Convergent Validity

The MEX: Descriptive Statistics and Internal Consistency

The means and standard deviations for the MEX barriers and benefits factors in each sample are shown in Table 4. The coefficients alpha [35] and omega [36] were also calculated for each factor in each sample and reflected good internal consistency across each sample and factor, with estimates ranging between 0.900 and 0.951 (see Table 4).

The MEX: Convergent Validity

The correlations between the MEX subscales and each of the pre-existing scales are shown in Table 5, along with means and standard deviations of each, and Fisher’s Z scores denoting comparative strengths of correlation coefficients between each variable and benefits and barriers, respectively.

In sample 1, the benefits scale was most strongly associated with satisfaction with life, and subjective ratings of physical activity and fitness; both correlation coefficients were significant and positive. The barriers score for sample 1 was most strongly associated with depression, anxiety, stress, self-deceptive enhancement, and impression management. Depression, anxiety and stress were each positively associated with the barriers scale, whilst self-deceptive enhancement and impression management were negatively associated with barriers to

Table 4 Means, standard deviations, minimum and maximum, and median scores of the MEX factors as summed total scores

	Sample 1	Sample 2	Sample 3
MEX-benefits			
Mean (SD)	46.8 (6.96)	46.1 (7.41)	45.6 (7.76)
Min/Max score	15–60	27–60	15–60
Median	48	45	45
Cronbach’s alpha	0.900	0.936	0.942
Omega	0.900	0.937	0.943
MEX-barriers			
Mean (SD)	35.8 (11.7)	35.3 (8.73)	35.8 (8.94)
Min/max score	15–56	15–54	15–57
Median	36	36	36
Cronbach’s alpha	0.951	0.906	0.910
Omega	0.951	0.910	0.916

Table 5 Convergent validity estimates for the MEX benefits and barriers subscales and existing scales

Scale	N	M (SD)	Benefits r	Barriers r	Fisher’s Z
Sample 1					
MEX-Benefits	492	46.8 (6.96)	–	–	–
MEX-Barriers	492	35.8 (11.7)	0.1*†	–	–
DASS-Depression	466	16.9 (12.6)	0.03	0.725***	12.314***
DASS-Anxiety	457	17.3 (13.7)	0.111*	0.703***	10.51***
DASS-Stress	462	16.9 (11.6)	0.093*	0.707***	10.881***
SWLS	482	22.8 (7.26)	0.271***	0.092*	2.719**
MET-mins/week	344	2972 (2102)	0.276***	0.164**	1.407
SPAF	480	13.2 (3.42)	0.377***	0.097*	4.363***
SDE	492	4.15 (0.782)	0.149**	–0.337***	7.462***
IM	492	4.21 (0.822)	–0.018	–0.254***	3.582***
Sample 2/3					
MEX-Benefits	605	45.8 (7.58)	–	–	–
MEX-Barriers	605	35.6 (8.83)	–0.454***	–	–
MASQ-AA	605	28.1 (10.1)	–0.033	0.306***	5.003***
MASQ-AD	605	97.5 (12.0)	–0.195***	0.466***	10.086***
MASQ-GDD	605	30.1 (11.1)	–0.174***	0.464***	9.721***
MASQ-GDA	605	23.2 (7.56)	–0.107**	0.394***	7.527***
MASQ-GDM	605	38.9 (11.9)	–0.154***	0.459***	9.337***
SPAF	605	10.9 (3.18)	0.437***	–0.606***	16.823***
LTSES	605	0.815 (0.99)	0.335***	–0.451***	11.998***

Correlations significant at $p < .001$ are emphasized in bold
 DASS Depression Anxiety Stress Scale, SWLS Satisfaction with Life Scale, MET Metabolic Equivalents of Physical Activity (minutes per week), SPAF Subjective Physical Activity and Fitness, LTSES Leisure-Time Sport and Exercise Score, SDE Self-Deceptive Enhancement, IM Impression Management, MASQ Mood, Anxiety, and Stress Questionnaire, AA Anxious Arousal, AD Anhedonic Depression, GDD General Distress: Depression, GDA General Distress: Anxiety, GDM = General Distress: Mixed

* $p < .05$, ** $p < .005$, *** $p < .001$

† Non-parametric relationship

exercise. Physical activity (PA) correlated with each scale with a similar magnitude, both with moderate positive associations.

In sample 2/3, the barriers score was significantly more associated with all variables than was the benefits score. The barriers score was positively correlated with all five MASQ-90 anxiety, depression and stress-related subscales. The barriers score was negatively associated with subjective ratings of physical activity and fitness, and with leisure-time exercise.

Barriers and Benefits Association

In both data sets, initial inspections of the association between the MEX barrier and benefit subscales were quantified using Pearson’s correlation. The two variables showed a moderate and linear negative correlation in sample 2/3 ($r = -0.454, p < 0.001$), but a small positive correlation in sample 1 ($r = 0.1, p = 0.03$). Upon further inspection, the relationship between barriers and benefits in sample 1 appeared to be nonlinear (see Fig. 1). A stepwise multiple regression model predicting benefits from the barrier values and the squared barrier values implied a quadratic relationship, $F(2, 489) = 34.77, p < 0.001, \text{adjusted } R^2 = 0.121$. Both the barrier values and the squared barrier values were statistically significant predictors of benefits in the stepwise regression model.

COVID-19 Impact on Activity and Mental Health

In sample 1, the impact of the COVID-19 pandemic on exercise behaviours was not significantly correlated with neither the MEX benefits ($r = -0.18, p = 0.078$), nor the MEX barriers ($r = -0.03, p = 0.759$). Similarly, the impact of the COVID-19 pandemic on mental health was not significantly correlated with neither the MEX benefits ($r = 0.09, p = 0.282$) nor the MEX barriers ($r = -0.06, p = 0.499$).

Discussion

Results from the present study support the psychometric validity and reliability of the 30-item Mental health-related barriers and benefits to EXercise (MEX) scale in generally healthy adults without severe mental illness. Additionally, the two 15-item subscales (benefits, and barriers) are each well supported psychometrically within

the present analysis. Both subscales have high item loadings and high internal consistency, whilst the scale as a whole showed excellent discriminant and convergent validity. The novel MEX scale fills an important gap in the literature regarding the quantification of depression and anxiety-related barriers and benefits to exercise engagement, which are well documented in qualitative research [4, 12], but are lesser known in quantitative research and not yet a key focus of any quantitative measurement of barriers and benefits of exercise.

There was a robust positive association in sample 1 among benefits and total physical activity (PA; i.e. a summation of weekly leisure-time exercise, as well as work, transport, or other non-leisure-time physical activity across an average week). In sample 2/3, there was also a robust positive association between the MEX benefits and leisure-time exercise. These results are important in showing the convergent validity of the MEX in terms of both overall PA, and in leisure-time physical activity, in particular. Similarly, barriers to exercise engagement were robustly positively associated with anxiety- and depression-like symptoms in both data sets and negatively associated with leisure-time exercise in sample 2/3. In both data sets, barriers were significantly more associated with symptoms of affective distress than were the benefits.

Given the number of overall correlations, only those significant at a threshold of $p < 0.001$ are considered in detail. In sample 1, results demonstrated a moderate positive correlation between benefits of exercise and satisfaction with life as predicted. Similarly, benefits of exercise were positively associated with total PA and subjective perception of fitness, suggesting that beliefs about benefits of exercise are related to both actual exercise engagement and the subjective estimates of one’s physical fitness and PA levels. Barriers, on the other hand, were strongly positively correlated with anxiety-like symptoms, depression-like symptoms, and self-reported levels of stress. This result implies that as symptoms of mental ill health increase, so too does one’s experience of mental health-related barriers to exercise engagement. The trend is particularly important to recognize as it further highlights the additional and unique barriers to exercise faced by otherwise generally healthy individuals who experience symptoms of anxiety and depression, as previously examined in qualitative research [4, 12]. To our knowledge, these results mark the first time that the relationships between mental health-related barriers to exercise and symptoms of anxiety, depression and stress have been quantified. Interestingly, there were moderate negative correlations between the barriers scale and both indicators of social desirability biases: impression management and self-deceptive enhancement. These correlations




Fig. 1 Raw barrier scores by raw benefit scores plotted with curvilinear prediction line

potentially imply that mental health-related barriers to exercise may be underreported by individuals with higher propensity for social desirability biases.

In sample 2/3, the mental health-related benefits to exercise were negatively correlated with symptoms related to anhedonic depression, symptoms related to general distress—depression, and general distress—mixed, as well as symptoms related to general distress—anxiety to a lesser extent. The results would seem to suggest a stronger relationship between benefits and depressive symptoms than benefits and anxious symptoms, though this was not observed in sample 1. Benefits were again positively correlated with the subjective assessment of physical fitness and PA, as well leisure-time exercise and sport engagement, providing important support for the scale's convergent validity in physical activity measurement and self-perception. The barriers scale was positively correlated with all measures of affective distress symptoms in sample 2/3, where the MASQ-90 was employed. Thus, for each of the five affective distress subscales, higher levels of symptoms corresponded with higher levels of perceived mental health barriers. Additionally, in sample 2/3, the barriers scale was negatively associated with leisure-time exercise, as well as subjective fitness and PA. Thus, higher levels of perceived barriers corresponded with lower levels of exercise, and lower levels of perceived fitness and activity in generally healthy adults without severe mental illness.

Some inconsistency between samples was observed. Firstly, the benefits scale correlated significantly with small and negative effects, with nearly all forms of distress in sample 2/3 except anxious arousal. However, in sample 1, the benefits scale showed only small, positive correlations with anxiety and stress and was not correlated with depression. This particular discrepancy may be due to the distributions of scores of the DASS-42 and the MASQ-90 in data sets 1 and 2, respectively. Whilst the DASS-42 subscales each demonstrated bimodal distributions in sample 1, the MASQ-90 subscales each demonstrated positively skewed distributions in sample 2/3. The benefits subscale scores were negatively skewed in both data sets 1 and 2. Thus, there may be a potential effect of the discrepancies in distributions of anxious and depressive-type symptoms on the reporting of benefits. Future research may seek to include more homogenous clinical samples for a clearer focus of the relationship between reported benefits and mental ill health symptoms.

Secondly, the barriers scale correlated significantly with moderate and negative effects, with both leisure-time physical activity and with subjective fitness in sample 2/3. However, in sample 1, the barriers scale correlated positively but weakly with self-reported metabolic equivalent minutes of PA, and with subjective fitness. Indeed, a

key difference arises in the measurement of leisure-time exercise specifically in sample 2/3, compared with using overall PA in sample 1. Given that the questions of the MEX scale ask for self-reported attitudes toward leisure-time exercise behaviours and not all physical activity and movement, it is reasonable to consider the leisure-time exercise scale, as used in sample 2/3, as a better counterpart to the MEX than the measurement of overall PA which was used in sample 1. In consideration of subjective physical activity, it's worthwhile to note that the distributions of scores differed between data sets. In sample 1, a negatively skewed distribution was observed for subjective physical activity, which coincided with a very low correlation between barriers and subjective activity ($r=0.097$). Accordingly, previous evidence also suggests a tendency for individuals who are more sedentary to overestimate PA on the IPAQ [22], the scale used in sample 1 only. In sample 2/3, an approximately normal distribution was observed and coincided with a moderate correlation between barriers and subjective activity ($r=-0.606$). Thus, whilst high levels of barriers may be prevalent in individuals who consider their activity and fitness to be low, there may still be a tendency of some individuals to experience higher than expected levels of barriers, despite self-reporting high levels of fitness and physical activity. Such a relationship may be moderated by additional factors, such as affective symptoms, psychological stress, and stages of change for exercise [37], which is a potential area for future exploration. It is worth noting that in some instances, higher levels of self-reported psychological stress has correlated with higher overall PA in young adults [37], but the exact relationship between the MEX barriers scale and differing types of PA beyond leisure-time exercise (e.g. occupational, housework, or transport PA) may still require further examination.

Finally, an additional inconsistency resulted from the relationship between benefits and barriers was inconsistent across data sets. In sample 2/3, a moderate negative linear relationship was found between MEX barrier and benefit scores. However, a moderate curvilinear association was found in sample 1, whereby low levels of barriers and high levels of barriers were both associated with high levels of benefits, but medium levels of barriers were associated with low levels of benefits. Thus, there appears to be potential variability amongst individuals with high levels of mental health barriers to exercise. Previous research has indicated that some, but not all, individuals may increase their overall physical activity in the face of high levels of psychological stress or mental health difficulty, choosing to use physical activity as a form of stress relief [37]. Indeed, the MEX includes stress relief as a theme within multiple items

of the benefits scale, potentially explaining why some individuals who report high levels of barriers may also report high levels of benefits. Nevertheless, the source of this variability amongst individuals who strongly endorse the barrier items will be an important focus in future research.

Limitations of the present work warrant consideration. Most importantly, the samples were limited to generally healthy adults without severe mental illness and largely with subclinical symptoms. Whilst the recruitment of generally healthy adults for scale validation was an important first step in validating the MEX, we recognize that the use of clinical samples for validation is crucial to future research with the MEX.

Secondly, comparison between data sets was made difficult by differences in measurement scales, especially for symptoms of depression, anxiety, and stress, and for leisure-time physical activity. Future research using the MEX in separate samples should therefore use consistent measures of depression, anxiety, and stress (i.e. the DASS-42), and leisure-time physical activity (i.e. the LTSES). Additionally, we did not include an objective measure of physical activity. It is important for future work to examine how items on the MEX relate to objective indicators of physical activity, given the typical low reliability found in self-report measures of activity. However, a differentiation may need to be made between leisure-time physical activity and total physical activity in objective measurement. Furthermore, future research should also examine the effects of mental health-related barriers and benefits on separable parameters of exercise prescriptions, including frequency, duration and intensity of exercise, as each parameter may have differing associations with the barriers and benefits scales [38], which may be important in personalization of exercise prescription.

It would also be important for future research to examine the potential effects of individual items or themes of the MEX (e.g. aversive sensations as barriers, or stress relief as a benefit). Due to the factor structure of the scale, whereby subfactors within the barrier and benefit factors could not be established in this study, conclusions about individual items or themes could not be reached. Future research could examine individual differences across MEX items, as certain items or themes may be of particular importance for individualization of treatment.

Finally, the sampling was limited to mostly Caucasian, English-speaking participants from western countries, with mostly subclinical depressive and anxious symptoms. Thus, ascertaining norms and factorial invariance for the MEX across diverse sociocultural samples, and in clinical samples, is an important consideration for future research.

Conclusions

We have developed a 30-item scale of mental health-related benefits and barriers to exercise, comprising 15 barriers and 15 benefits. The aim of this research was to produce a quantitative measurement of the bidirectional relationship between transdiagnostic symptoms of mental health disorders and physical exercise, which can be used at both the individual level in clinical settings, and at broader levels for research. The present research provides evidence that scores on the MEX are valid and reliable in generally healthy adults without severe mental illness, as highlighted by the MEX subscales meeting excellent standards of internal consistency, and factorial, convergent, and discriminant validity. We believe that due to these properties, as well as the brevity and breadth of the scale, the MEX will be a useful measurement tool in both clinical and research settings for the mental health-related barriers and benefits to exercise.

Abbreviations

MEX	The Mental health-related barriers and benefits to EXercise scale
DASS-42	Depression Anxiety Stress Scale
SWLS	Satisfaction with Life Scale
METS	Metabolic Equivalents of Physical Activity
SPAF	Subjective Physical Activity and Fitness
LTSES	Leisure-Time Sport and Exercise Score
SDE	Self-Deceptive Enhancement
IM	Impression Management
MASQ	Mood, Anxiety, and Stress Questionnaire
MASQ-AA	Anxious Arousal
MASQ-AD	Anhedonic Depression
MASQ-GDD	General Distress: Depression
MASQ-GDA	General Distress: Anxiety
MASQ-GDM	General Distress: Mixed
PA	Physical Activity
IRV	Intra-individual response variability
EFA	Exploratory Factor Analysis
CFA	Confirmatory Factor Analysis
TLI	Tucker–Lewis Index
CFI	Comparative fit index
SRMR	Standardized Root-Mean-Squared Residual
RMSEA	Root-mean-square error of approximation

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s40798-023-00555-x>.

Additional file 1: Supplementary demographic information, and the data cleaning procedure for each of the samples analysed.

Acknowledgements

The authors would like to acknowledge Professor Željko Pedišić (Victoria University, Australia) for his advice on material choice and methodology

Author contributions

M.L.C. contributed to idea conception, item and scale development, data collection, data analysis, and wrote the manuscript. S.C.B. assisted with data analysis and edited the manuscript. M.C.P. edited the manuscript. N.T.V.D. assisted with item and scale development, assisted with data collection,

assisted with data analysis, and edited the manuscript. All authors read and approved the final manuscript.

Funding

This research was financially supported by the Melbourne School of Psychological Sciences at the University of Melbourne.

Availability of Data and Materials

Data are available online: <https://osf.io/dm3kz/>. The MEX scale is available online: <https://osf.io/gnqfr/wiki/home/>. Analysis scripts are available upon request to the corresponding author. The availability of other materials as described in the Methods section will be at the discretion of the copyright holders per each item.

Declarations

Ethics Approval and Consent to Participate

Ethics approved by the University of Melbourne Human Research Ethics Committee [HREC # 2056662]. All participants were provided plain language statements and consent forms prior to participating in the research. Only participants who confirmed their informed consent could continue to participate in the research tasks. The present research conforms to the standards of ethics outlined in the Declaration of Helsinki.

Consent for Publication

Consent form as above notified all participants of our intent to publish.

Competing Interests

We declare no conflict of interest nor competing interests in the authorship or funding of this paper.

Received: 11 August 2022 Accepted: 24 January 2023

Published online: 23 February 2023

References

- WHO. Depression and Other Common Mental Disorders: Global Health Estimates. World Health Organization, Geneva, 2017. Accessed: 29 April 2022. Available: <https://www.who.int/news-room/fact-sheets/detail/mental-disorders>
- Rebar AL, Stanton R, Geard D, Short C, Duncan MJ, Vandelanotte C. A meta-meta-analysis of the effect of physical activity on depression and anxiety in non-clinical adult populations. *Health Psychol Rev*. 2015;9(3):366–78. <https://doi.org/10.1080/17437199.2015.1022901>.
- Glowacki K, Duncan MJ, Gainforth H, Faulkner G. Barriers and facilitators to physical activity and exercise among adults with depression: a scoping review. *Ment Health Phys Act*. 2017;13:108–19. <https://doi.org/10.1016/j.mhpa.2017.10.001>.
- Searle A, Calnan M, Lewis G, Campbell J, Taylor A, Turner K. Patients' views of physical activity as treatment for depression: a qualitative study. *Br J Gen Pract*. 2011;61(585):e149–56. <https://doi.org/10.3399/bjgp11X567054>.
- Cooney GM, et al. Exercise for depression. *Cochrane Database of Systematic Reviews*, no. 9, 2013. <https://doi.org/10.1002/14651858.CD004366.pub6>.
- White RL, Babic MJ, Parker PD, Lubans DR, Astell-Burt T, Lonsdale C. Domain-specific physical activity and mental health: a meta-analysis. *Am J Prev Med*. 2017;52(5):653–66. <https://doi.org/10.1016/j.amepre.2016.12.008>.
- Sallis JF. Age-related decline in physical activity: a synthesis of human and animal studies. *Med Sci Sports Exerc*. 2000;32(9):1598–600.
- Solmi M, et al. Age at onset of mental disorders worldwide: large-scale meta-analysis of 192 epidemiological studies. *Mol Psychiatry*, 27(1), Art. no. 1, 2022; <https://doi.org/10.1038/s41380-021-01161-7>.
- Sechrist KR, Walker SN, Pender NJ. Development and psychometric evaluation of the exercise benefits/barriers scale. *Res Nurs Health*. 1987;10(6):357–65. <https://doi.org/10.1002/nur.4770100603>.
- Burton NW, Oldenburg B, Sallis JF, Turrell G. Measuring psychological, social, and environmental influences on leisure-time physical activity among adults. *Aust N Z J Public Health*. 2007;31(1):36–43.
- Chapman JJ, Fraser SJ, Brown WJ, Burton NW. Physical activity preferences, motivators, barriers and attitudes of adults with mental illness. *J Ment Health*. 2016;25(5):448–54. <https://doi.org/10.3109/09638237.2016.1167847>.
- Mason JE, Faller YN, LeBouthillier DM, Asmundson GJG. Exercise anxiety: a qualitative analysis of the barriers, facilitators, and psychological processes underlying exercise participation for people with anxiety-related disorders. *Ment Health Phys Act*. 2019;16:128–39. <https://doi.org/10.1016/j.mhpa.2018.11.003>.
- Seime RJ, Vickers KS. The challenges of treating depression with exercise: from evidence to practice. *Clin Psychol Sci Pract*. 2006;13(2):194–7.
- Mikkelsen K, Stojanovska L, Polenakovic M, Bosevski M, Apostolopoulos V. Exercise and mental health. *Maturitas*. 2017;106:48–56. <https://doi.org/10.1016/j.maturitas.2017.09.003>.
- Mundfrom DJ, Shaw DG, Ke TL. Minimum sample size recommendations for conducting factor analyses. *Int J Test*. 2005;5(2):159–68. https://doi.org/10.1207/s15327574ijt0502_4.
- Brown SA, Huber D, Bergman A. A perceived benefits and barriers scale for strenuous physical activity in college students. *Am J Health Promot*. 2006;21(2):137–40. <https://doi.org/10.4278/0890-1171-21.2.137>.
- Busch AM, Ciccolo JT, Puspitasari AJ, Nosrat S, Whitworth JW, Stults-Kolehmainen MA. Preferences for exercise as a treatment for depression. *Ment Health Phys Act*. 2016;10:68–72. <https://doi.org/10.1016/j.mhpa.2015.12.004>.
- Firth J, Rosenbaum S, Stubbs B, Gorczynski P, Yung AR, Vancampfort D. Motivating factors and barriers towards exercise in severe mental illness: a systematic review and meta-analysis. *Psychol Med*. 2016;46(14):2869–81. <https://doi.org/10.1017/S0033291716001732>.
- Ussher M, Stanbury L, Cheeseman V, Faulkner G. Physical activity preferences and perceived barriers to activity among persons with severe mental illness in the United Kingdom. *Psychiatr Serv*. 2007;58(3):405–8.
- Lovibond SH, Lovibond PF. Manual for the Depression Anxiety Stress Scales, 2nd ed. Psychology Foundation, 1995.
- Booth M. Assessment of physical activity: an international perspective. *Res Q Exerc Sport*. 2000;71(sup2):114–20. <https://doi.org/10.1080/02701367.2000.11082794>.
- Shook R, et al. Subjective estimation of physical activity using the IPAQ varies by fitness level. *J Phys Activity Health*, 13, 2015; <https://doi.org/10.1123/jpah.2014-0543>.
- Diener E, Emmons RA. The satisfaction with life scale. *J Personality Assessment*, p. 5, 1985.
- Hart CM, Ritchie TD, Hepper EG, Gebauer JE. The balanced inventory of desirable responding short form (BIDR-16). *SAGE Open*. 2015. <https://doi.org/10.1177/2158244015621113>.
- Lang C, Brand S, Colledge F, Ludyga S, Pühse U, Gerber M. Adolescents' personal beliefs about sufficient physical activity are more closely related to sleep and psychological functioning than self-reported physical activity: a prospective study. *J Sport Health Sci*. 2019;8(3):280–8. <https://doi.org/10.1016/j.jsjshs.2018.03.002>.
- Watson D, Clark LA, Weber K, Assenheimer JS, Strauss ME, McCormick RA. Testing a tripartite model: II. Exploring the symptom structure of anxiety and depression in student, adult, and patient samples. *J Abnorm Psychol*. 1995;104(1):15–25. <https://doi.org/10.1037//0021-843x.104.1.15>.
- Baecke JA, Burema J, Frijters JE. A short questionnaire for the measurement of habitual physical activity in epidemiological studies. *Am J Clin Nutr*. 1982;36(5):936–42. <https://doi.org/10.1093/ajcn/36.5.936>.
- Meade AW, Craig SB. Identifying careless responses in survey data. *Psychol Methods* 17(3):437.
- Dunn AM, Heggstad ED, Shanock LR, Theilgard N. Intra-individual response variability as an indicator of insufficient effort responding: comparison to other indicators and relationships with individual differences. *J Bus Psychol*. 2018;33(1):105–21. <https://doi.org/10.1007/s10869-016-9479-0>.
- Hu L, Bentler PM. Cutoff criteria for fit indexes in covariance structure analysis: conventional criteria versus new alternatives. *Null*. 1999;6(1):1–55. <https://doi.org/10.1080/10705519909540118>.
- Brown TA. Confirmatory factor analysis for applied research. Guilford publications; 2015.

32. Kline P. An Easy guide to factor analysis. London: Routledge, 2014; <https://doi.org/10.4324/9781315788135>.
33. Dunn OJ, Clark V. Correlation coefficients measured on the same individuals. *J Am Stat Assoc.* 1969;64(325):366–77.
34. Drasgow F, Lissak RI. Modified parallel analysis: a procedure for examining the latent dimensionality of dichotomously scored item responses. *J Appl Psychol.* 1983;68(3):363–73. <https://doi.org/10.1037/0021-9010.68.3.363>.
35. Cronbach LJ. Essentials of psychological testing; 1970.
36. Bollen KA. Structural equations with latent variables, vol. 210. Wiley, 1989.
37. Stults-Kolehmainen MA, Sinha R. The effects of stress on physical activity and exercise. *Sports Med.* 2014;44(1):81–121. <https://doi.org/10.1007/s40279-013-0090-5>.
38. Lutz RS, Stults-Kolehmainen MA, Bartholomew JB. Exercise caution when stressed: stages of change and the stress–exercise participation relationship. *Psychol Sport Exerc.* 2010;11:560–7. <https://doi.org/10.1016/j.psychsport.2010.06.005>.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Submit your manuscript to a SpringerOpen[®] journal and benefit from:

- ▶ Convenient online submission
- ▶ Rigorous peer review
- ▶ Open access: articles freely available online
- ▶ High visibility within the field
- ▶ Retaining the copyright to your article

Submit your next manuscript at ▶ [springeropen.com](https://www.springeropen.com)
